

Active Motif Antibodies

The table below provides details on all of Active Motif's antibodies and was created on May 27, 2011. For an up-to-date list and for more comprehensive information on each antibody, including data and downloadable Technical Data Sheets, please visit our website at www.activemotif.com/abs. To see our Fluorescent and HRP Secondary Antibody Conjugates, please go to the last page of this antibody list or visit us at www.activemotif.com/secondary.

	Antibody	Immunogen	Reactivity	Applications	Format	Catalog No.
NEW	5-Hydroxymethylcytidine antibody (mAb) (Clone 59.1)	Raised against 5-Hydroxymethylcytidine conjugated to KLH.	H, M, WR	DB, MeDIP	100 µg	39999
					10 µg	40000
	5-Hydroxymethylcytidine antibody (pAb)	Raised against 5-Hydroxymethylcytidine conjugated to KLH.	H, M, WR	DB, IF, IHC, MeDIP	100 µl	39769
					10 µl	39770
	5-Hydroxymethylcytidine antibody (pAb)	Raised against 5-Hydroxymethylcytidine conjugated to KLH.	H, M, WR	DB, MeDIP	100 µg	39791
					10 µg	39792
	5-Methylcytidine antibody (mAb) (Clone 33D3)	Recognizes the modified 5-methylcytidine base in plant and vertebrate DNA.	WR	FC, IF, IHC, IP	50 µg	39649
	Bridging Antibody for Mouse IgG	Raised against purified mouse IgG.	M	ChIP, IP	500 µg	53017
	A20 antibody (mAb)	Raised against full-length human A20.	H	IP, WB	100 µg	40901
	Acf1 antibody (pAb)	Raised against the recombinant, C-terminal half of <i>Drosophila</i> Acf1.	Dr	IF, IP, WB	100 µl	39721
					10 µl	39722
	Ago1/2/3 antibody (mAb) (Clone 2A8)	Raised against a recombinant protein with amino acids 47-879 of human Ago2.	H	ChIP, IF, IHC, IP, WB	100 µg	39937
					10 µg	39938
NEW	Ago1 antibody (mAb) (Clone 4B8)	Raised against a peptide containing the N-terminal region of human Ago1.	H	IP, WB	100 µg	61071
					10 µg	61072
	Ago2 antibody (mAb) (Clone 1B1-E2H5)	Raised against a recombinant protein with amino acids 1-148 of human Ago2.	H	IF, IP, WB	100 µg	39853
					10 µg	39854
	Ago3 antibody (mAb) (Clone 4B1-F6)	Raised against a recombinant protein with amino acids 1-175 of human Ago3.	H, M	IF, IP	100 µg	39787
					10 µg	39788
	Ago4 antibody (mAb) (Clone 1B7-G11)	Raised against a recombinant protein with amino acids 1-164 of human Ago4.	H	IP, WB	100 µg	39855
					10 µg	39856
	Ago4 antibody (pAb)	Raised against a peptide containing the N-terminal region of human Ago4.	H	WB	100 µl	39813
					10 µl	39814
	AIB1 / SRC-3 antibody (pAb)	Raised against a peptide derived from human AIB1 / SRC-3.	H	ChIP, WB	100 µl	39797
					10 µl	39798
	AIB3 antibody (pAb)	Raised against a recombinant protein with amino acids 2-146 of human AIB3.	H, M	IHC, IP, WB	100 µl	39835
					10 µl	39836
	AID antibody (mAb) (Clone 328.8b)	Raised against amino acids 178-198 of human AID.	H, M	IF, IP, WB	100 µg	39885
					10 µg	39886
	Aiolos antibody (pAb)	Raised against a GST-tagged, C-terminal half of mouse Aiolos.	M	IF, IP, WB	200 µl	39293
					10 µl	39294

Reactivity: B = Bovine; BY = Budding Yeast; Ce = *C. elegans*; Dr = *Drosophila*; FY = Fission Yeast; Hm = Hamster; H = Human; Mk = Monkey; M = Mouse; NS = Not Species Specific; Rb = Rabbit; R = Rat; WR = Wide range predicted
Applications: ChIP = Chromatin IP; DB = Dot Blot; FC = Flow Cytometry; ICC = Immunocytochemistry; IF = Immunofluorescence; IHC = Immunohistochemistry; IP = Immunoprecipitation; MeDIP = Methyl-DNA IP; WB = Western Blot

Antibody	Immunogen	Reactivity	Applications	Format	Catalog No.
Aiolos antibody (mAb) (Clone 8B2)	Raised against the N-terminal half of mouse Aiolos.	M	ChIP, WB	100 µg 10 µg	39657 39658
AKT1 phospho Ser473 antibody (mAb)	Raised against a peptide with phospho-serine 473 of human AKT1.	H, M	WB	100 µg	40902
alpha Tubulin antibody (mAb) (Clone 5-B-1-2)	Raised against total protein purified from sea urchin sperm axonemes.	H, M, WR	IF, WB	200 µl 10 µl	39527 39528
AML-1/Runx1 antibody (pAb)	Raised against amino acids 231-245 of human AML-1/Runx1.	H, M	WB	100 µl	39000
AML-2/Runx3 antibody (pAb)	Raised against full-length human AML-2/Runx3.	H	EMSA, WB	100 µl	39301
AML-3/Runx2 antibody (pAb)	Raised against amino acids 333-348 of human AML-3/Runx2.	H	EMSA	100 µl	39302
Androgen Receptor antibody (pAb)	Raised against a peptide of the N-terminal region of human Androgen Receptor.	H	WB	100 µl 10 µl	39781 39782
AP-2 antibody (pAb)	Raised against amino acids 120-134 of human AP-2.	H	WB	100 µg	39001
APE antibody (pAb)	Peptide mixture of amino acids 36-52 and 216-232 of human APE.	H	WB	100 µg	39200
APEXL2 antibody (pAb)	Raised against a peptide corresponding to human APEXL2.	H	WB	200 µl	39201
APOBEC2 antibody (pAb)	Raised against a peptide of the N-terminal region of human APOBEC2.	H	WB	100 µl 10 µl	39985 39986
Dimethyl-Arginine, asymmetric antibody (pAb)	Raised against a peptide with dimethyl-arginine (asymmetric).	NS	IF, WB	200 µl 10 µl	39231 39232
ASH2 antibody (pAb)	Raised against full-length recombinant human ASH2 protein.	H, WR	IF, IP, WB	200 µl 10 µl	39099 39100
ATF-2 antibody (pAb)	Raised against amino acids 13-27 of human ATF-2.	H	WB	100 µg	40977
ATF-2 phospho Thr71 antibody (mAb)	Raised against a peptide with phospho-threonine 71 of human ATF-2.	H	WB	100 µg	40975
ATF-6 antibody (mAb)	Raised against a partial protein of amino acids 1-273 of human ATF-6.	H	IF, WB	100 µg	40962
ATM phospho Ser1981 antibody (mAb) (Clone 10H11.E.12)	Raised against a peptide with phospho-serine 1981 of human ATM.	H, M	IF	200 µl 10 µl	39529 39530
NEW ATRX antibody (pAb)	Raised against a peptide with the C-terminus of human ATRX.	H	WB	200 µl 10 µl	61049 61050
Aurora B antibody (pAb)	Raised against a peptide with the C-terminus of human Aurora B.	H, M, WR	IF, WB	200 µl 10 µl	39261 39262
Bcl3 antibody (pAb)	Raised against a C-terminal peptide of human Bcl3.	H	WB	100 µl 10 µl	39761 39762
Bcl10 antibody (mAb) (Clone 151)	Raised against full-length recombinant human Bcl10 protein.	H	WB	200 µg 10 µg	39393 39394
NEW BCL11A / CTIP1 antibody (pAb)	Raised against a peptide of the N-terminal region of human BCL11A / CTIP1.	H	WB	100 µl 10 µl	61051 61052
NEW BLIMP1 / PRDM1 antibody (pAb)	Raised against a peptide of the N-terminal half of human BLIMP1 / PRDM1.	H	WB	100 µl 10 µl	61053 61054

Reactivity: B = Bovine; BY = Budding Yeast; Ce = *C. elegans*; Dr = *Drosophila*; FY = Fission Yeast; Hm = Hamster; H = Human; Mk = Monkey; M = Mouse; NS = Not Species Specific; Rb = Rabbit; R = Rat; WR = Wide range predicted
Applications: ChIP = Chromatin IP; DB = Dot Blot; FC = Flow Cytometry; ICC = Immunocytochemistry; IF = Immunofluorescence; IHC = Immunohistochemistry; IP = Immunoprecipitation; MeDIP = Methyl-DNA IP; WB = Western Blot

	Antibody	Immunogen	Reactivity	Applications	Format	Catalog No.
NEW	BLM antibody (pAb)	Raised against a peptide of the N-terminal region of human BLM.	H	WB	100 µl 10 µl	61055 61056
	BMI-1 antibody (mAb) (Clone AF27)	Raised against full-length recombinant mouse BMI-1 protein.	H, M	ChIP, IP	100 µg 10 µg	39993 39994
NEW	BMI-1 antibody (mAb) (Clone DC9)	Raised against full-length recombinant mouse BMI-1 protein.	H, M	IF, WB	100 µg 10 µg	61089 61090
	Boris / CTCFL antibody (pAb)	Raised against a peptide of the N-terminal region of human Boris.	H	WB	100 µl 10 µl	39851 39852
NEW	BRD4 antibody (pAb)	Raised against a C-terminal peptide of human BRD4.	H	WB	100 µl 10 µl	39909 39910
	BRD8 / SMAP2 antibody (pAb)	Raised against a peptide of the N-terminal region of human BRD8 / SMAP2.	H	WB	100 µl 10 µl	61007 61008
NEW	BRG-1 antibody (mAb) (Clone 3G4)	Raised against a recombinant protein containing amino acids 214-279 of mouse BRG-1.	H	IF, WB	100 µg 10 µg	39807 39808
	BRM antibody (mAb) (Clone 1H7A10)	Raised against a recombinant protein containing amino acids 48-214 of mouse BRM.	H	ChIP, IF, WB	100 µg 10 µg	39805 39806
NEW	Bub1 antibody (mAb) (Clone 145H)	Raised against amino acids 281-419 of human Bub1.	H, M	IF, WB	100 µg 10 µg	39735 39736
	BubR1 antibody (mAb) (Clone 8G1)	Raised against amino acids 1-350 of human BubR1.	H, M, WR	IF, WB	100 µg 10 µg	39747 39748
NEW	c-Fos antibody (pAb)	Raised against amino acids 358-372 of human c-Fos.	H, M	WB	100 µl	39008
	c-Fos antibody (pAb)	Raised against amino acids 358-372 of human c-Fos.	H, M	EMSA	100 µl	39308
NEW	c-Jun antibody (pAb)	Raised against amino acids 20-37 of human c-Jun.	H, M	WB	100 µl	39009
	c-Jun antibody (pAb)	Raised against amino acids 20-37 of human c-Jun.	H, M	ChIP, EMSA, IF	100 µg	39309
NEW	c-Myc antibody (mAb) (Clone 6A10)	Raised against a recombinant protein containing amino acids 1-262 of human c-Myc.	H	EMSA, IP, WB	100 µg 10 µg	61075 61076
	c-Myc antibody (mAb) (61 kDa form)	Raised against amino acids 408-420 of human c-Myc; 61 kDa form.	H	WB	100 µl	39502
NEW	c-Myc antibody (pAb) (65 kDa form)	Raised against amino acids 9-22 of human c-Myc; detects 65 kDa form.	H, Mk, M	WB	100 µl	39012
	c-Rel antibody (pAb)	Raised against amino acids 507-524 of human c-Rel.	H	EMSA, WB	100 µg	39311
NEW	C/EBPα antibody (pAb)	Raised against amino acids 1-14 of human C/EBPα.	H, R	ChIP, EMSA, IF, WB	100 µg	39306
	C/EBPβ antibody (pAb)	Raised against amino acids 228-242 of human C/EBPβ.	H, R	EMSA, WB	100 µl	39307
NEW	CAF-1 p60 antibody (mAb) (Clone 24)	Raised against full-length CAF-1 protein purified from 293 cells.	H	IF, IP, WB	100 µg 10 µg	39995 39996
	CAF-1 p150 antibody (mAb) (Clone 1)	Raised against full-length CAF-1 protein purified from 293 cells.	H	IF, IP, WB	100 µg 10 µg	39997 39998
NEW	CARM1 antibody (pAb)	Raised against a peptide to the C-terminus of mouse CARM1.	H, M, R	WB	200 µl 10 µl	39251 39252

Reactivity: B = Bovine; BY = Budding Yeast; Ce = *C. elegans*; Dr = *Drosophila*; FY = Fission Yeast; Hm = Hamster; H = Human; Mk = Monkey; M = Mouse; NS = Not Species Specific; Rb = Rabbit; R = Rat; WR = Wide range predicted
Applications: ChIP = Chromatin IP; DB = Dot Blot; FC = Flow Cytometry; ICC = Immunocytochemistry; IF = Immunofluorescence; IHC = Immunohistochemistry; IP = Immunoprecipitation; MeDIP = Methyl-DNA IP; WB = Western Blot

Antibody	Immunogen	Reactivity	Applications	Format	Catalog No.
Caspase-3 antibody (mAb)	Raised against 6xHis-tagged, full-length human Caspase-3 protein.	H, M, R	WB	100 µg	40924
Caspase-7 antibody (mAb)	Raised against a recombinant human Caspase-7 protein.	H	WB	100 µg	40929
Caspase-8 antibody (mAb)	Raised against amino acids 217-222 of human Caspase-8.	H	FC, IHC, WB	100 µg	40930
Caspase-9 antibody (pAb) (Pro-form)	Raised against amino acids 299 -318 of human Pro-Caspase-9.	H	WB	100 µg	40931
Caspase-14 antibody (mAb)	Raised against amino acids 2-18 of mouse Caspase-14.	H, M	WB	100 µg	40932
NEW CBF-1 antibody (mAb) (Clone 7A11)	Raised against full-length recombinant human CBF-1.	H	WB	100 µg 10 µg	61073 61074
NEW CDT1 antibody (mAb)	Raised against a recombinant protein containing amino acids 177-557 of mouse CDT1.	H	IHC, IP, WB	100 µg 10 µg	61039 61040
CENP-E antibody (mAb) (Clone 1H12)	Raised against full-length human CENP-E.	H, M	IF	200 µg 10 µg	39619 39620
CGBP antibody (pAb)	Peptide mixture of amino acids 73-90 and 641-656 of human CGBP.	H, M	WB	200 µl	39203
CHD1 antibody (pAb)	Raised against amino acids 251-467 of human CHD1.	H	ChIP, WB	100 µl 10 µl	39729 39730
CHD1 antibody (mAb) (Clone 2F11)	Raised against a peptide to the C-terminus of mouse CHD1.	H, M	IF, WB	100 µg 10 µg	39989 39990
CHD2 antibody (pAb)	Raised against a peptide to the C-terminus of human CHD2.	H, WR	WB	200 µl 10 µl	39363 39364
CID antibody (pAb)	Raised against recombinant CID protein from <i>Drosophila</i> .	Dr	WB	100 µl 10 µl	39713 39714
CID antibody (pAb)	Raised against recombinant CID protein from <i>Drosophila</i> .	Dr	IF	100 µl 10 µl	39719 39720
Copine I antibody (pAb)	Raised against amino acids 519-532 of human Copine I.	H	WB	200 µl 10 µl	39101 39102
CoREST antibody (pAb)	Raised against a peptide to the C-terminal region of human CoREST.	H	WB	100 µl 10 µl	39955 39956
CREB antibody (mAb)	Raised against a peptide corresponding to human CREB.	H	WB	100 µg	40961
CREB-1 antibody (pAb)	Raised against amino acids 6-23 of human CREB-1.	H, R	WB	100 µl	39013
CTBP antibody (pAb)	Raised against a peptide to the C-terminal region of human CTBP.	H	WB	100 µl 10 µl	39847 39848
CTCF antibody (mAb)	Raised against human CTCF expressed in <i>Pichia pastoris</i> .	H, M	ChIP, WB	200 µg 10 µg	39621 39622
cyclin A antibody (mAb)	Raised against full-length human Cyclin A.	H	WB	100 µl	39503
cyclin A antibody (mAb)	Raised against full-length human Cyclin A.	H	EMSA	17 rxns	39600
cyclin B1 antibody (mAb)	Raised against a recombinant protein to human Cyclin B1.	H	WB	100 µl	39504
Cyk-4 phospho Ser170 antibody (pAb)	Raised against a peptide with phospho-serine 170 of human Cyk-4.	H	IF, WB	200 µl 10 µl	39265 39266

Reactivity: B = Bovine; BY = Budding Yeast; Ce = *C. elegans*; Dr = *Drosophila*; FY = Fission Yeast; Hm = Hamster; H = Human; Mk = Monkey; M = Mouse; NS = Not Species Specific; Rb = Rabbit; R = Rat; WR = Wide range predicted
Applications: ChIP = Chromatin IP; DB = Dot Blot; FC = Flow Cytometry; ICC = Immunocytochemistry; IF = Immunofluorescence; IHC = Immunohistochemistry; IP = Immunoprecipitation; MeDIP = Methyl-DNA IP; WB = Western Blot

Antibody	Immunogen	Reactivity	Applications	Format	Catalog No.
D4-GDI antibody (mAb)	Raised against the Fas-induced cleavage site of human D4-GDI.	H	IP, WB	100 µg	40941
DAP-3 antibody (mAb)	Raised against amino acids 51-66 of human DAP-3.	H, M	WB	100 µg	40963
DAP-5 antibody (mAb)	Raised against amino acids 796-812 of human DAP-5.	H, M	WB	100 µg	40964
DAX-1 / NR0B1 antibody (mAb) (Clone 1DA-2F4)	Raised against a peptide to amino acids 135-166 of human DAX-1 / NR0B1.	H, M, R	ELISA, ICC, IF, IHC, IP, WB	100 µg 10 µg	39983 39984
DcR3 antibody (mAb)	Raised against full-length human DcR3.	H	WB	100 µg	40933
Dicer antibody (mAb) (Clone NI67/7)	Raised against a recombinant protein containing amino acids 1638-1899 of mouse Dicer.	H, M	WB	100 µg 10 µg	39817 39818
DNMT1 antibody (mAb)	Raised against amino acids 637-651 of human DNMT1.	H, M	ChIP, IHC, IP, WB	100 µg	39204
DNMT1 antibody (pAb)	Raised against a recombinant protein containing amino acids 1-110 of human DNMT1.	H	WB	100 µl 10 µl	39905 39906
DNMT2 antibody (pAb)	Raised against amino acids 39-53 and 361-376 of mouse DNMT2.	H, M	WB	100 µg	39205
DNMT3A antibody (mAb)	Raised against full-length mouse DNMT3A.	H, M	ChIP, IF, IHC, WB	100 µg	39206
DNMT3A antibody (pAb)	Raised against a recombinant protein containing amino acids 1-247 of mouse DNMT3A.	H	WB	100 µl 10 µl	39897 39898
DNMT3B antibody (mAb)	Raised against full-length mouse DNMT3B.	H, M	ChIP, IF, IP, WB	100 µg	39207
DNMT3B antibody (pAb)	Raised against a recombinant protein containing amino acids 1-218 of mouse DNMT3B.	H	WB	100 µl 10 µl	39899 39900
DNMT3L antibody (pAb)	Raised against full-length recombinant human DNMT3L.	H	WB	100 µl 10 µl	39907 39908
DOT1L antibody (pAb)	Raised against a peptide to the C-terminal region of human DOT1L.	H	WB	100 µl 10 µl	39953 39954
DR4 antibody (mAb)	Raised against amino acids 1-20 of human DR4.	H	FC, WB	100 µg	40934
DR6 antibody (pAb)	Raised against amino acid residues 371-655 of human DR6.	H	WB	100 µg	40927
DRAK1 antibody (pAb)	Raised against amino acids 5 to 9 of human DRAK1.	H, M	WB	100 µg	40923
DRAK2 antibody (pAb)	Raised against amino acids 351-365 of human DRAK2.	H	WB	100 µg	40926
Drosha antibody (pAb)	Raised against a peptide containing the C-terminal region of human Drosha.	H	WB	100 µl 10 µl	39783 39784
E1A antibody (mAb) (Clone M73)	Raised against full-length recombinant Adenovirus 2 E1A.	Cells infected with Adenovirus	IF, IP, WB	100 µg 10 µg	39771 39772
E2F-1 antibody (mAb)	Raised against full-length human E2F-1.	H	EMSA	17 rxns	39601
E2F-1 antibody (pAb)	Raised against amino acids 309-323 of human E2F-1.	H, M	ChIP, EMSA	17 rxns	39313
E2F-3 antibody (mAb)	Raised against full-length human E2F-3.	H	WB	100 µl	39507
E2F-4 antibody (mAb)	Raised against full-length human E2F-4.	H	EMSA	17 rxns	39603
E2F-6 antibody (mAb)	Raised against full-length human E2F-6.	H	ChIP, WB	100 µl	39509
EGF Receptor phospho Tyr992 antibody (pAb)	Raised against a peptide with phospho-tyrosine 992 of human EGFR.	H	WB	200 µl	39094

Reactivity: B = Bovine; BY = Budding Yeast; Ce = *C. elegans*; Dr = *Drosophila*; FY = Fission Yeast; Hm = Hamster; H = Human; Mk = Monkey; M = Mouse; NS = Not Species Specific; Rb = Rabbit; R = Rat; WR = Wide range predicted
Applications: ChIP = Chromatin IP; DB = Dot Blot; FC = Flow Cytometry; ICC = Immunocytochemistry; IF = Immunofluorescence; IHC = Immunohistochemistry; IP = Immunoprecipitation; MeDIP = Methyl-DNA IP; WB = Western Blot

Antibody	Immunogen	Reactivity	Applications	Format	Catalog No.
Elf-1 antibody (pAb)	Raised against amino acids 89-102 of human Elf-1.	H	WB	100 µl	39019
ELP3 antibody (pAb)	Raised against a peptide to the N-terminal region of human ELP3.	H	WB	100 µl 10 µl	39949 39950
NEW Estrogen Receptor α antibody (mAb) (Clone F3)	Raised against a peptide containing the C-terminal region of human ER α .	H	ChIP, EMSA, IF, IP, WB	100 µg 10 µg	61035 61036
Estrogen Receptor α antibody (pAb)	Raised against amino acids 145-159 of human ER α .	H	IF, IHC, WB	100 µl	39021
Estrogen Receptor β antibody (mAb) (Clone CO1531)	Raised against amino acids 212-477 of human ER β .	H	ICC, IF, IP, WB	100 µg 10 µg	39767 39768
NEW ERR- α antibody (pAb)	Raised against a peptide containing the C-terminal region of human ERR- α .	H	WB	100 µl 10 µl	61047 61048
Ets-1 antibody (pAb)	Raised against the C-terminal half of recombinant mouse Ets-1.	H	WB	200 µl 10 µl	39580 39581
EuHMT1 antibody (pAb)	Raised against a peptide to the N-terminal region of human EuHMT1.	H	WB	100 µg 10 µg	39777 39778
EZH2 antibody (mAb) (Clone AC22)	Raised against a recombinant protein containing amino acids 353-451 of human EZH2.	H	ChIP, IF, IP	100 µg 10 µg	39875 39876
EZH2 antibody (pAb)	Raised against a recombinant protein containing amino acids 1-370 of mouse EZH2.	H, M	ChIP	100 µl 10 µl	39901 39902
EZH2 antibody (pAb)	Raised against a recombinant protein containing amino acids 1-370 of mouse EZH2.	H	WB	100 µl 10 µl	39933 39934
NEW FancD2 antibody (pAb)	Raised against a peptide within the internal region of human FancD2.	H	WB	100 µl 10 µl	61057 61058
FosB antibody (mAb)	Raised against amino acids 2-14 of human FosB.	H	WB	100 µg	40960
FosB antibody (pAb)	Raised against amino acids 2-14 of human FosB.	H, M	EMSA	17 rxns	39318
FosB antibody (pAb)	Raised against a peptide corresponding to human FosB.	H	WB	100 µg	40959
FOXO1/FKHR antibody (pAb)	Raised against a peptide to the C-terminal region of human FOXO1/FKHR.	H	WB	200 µl 10 µl	39629 39630
Fra-2 antibody (pAb)	Raised against amino acids 289-305 of human Fra-2.	H, M	WB	100 µl	39023
GAGA Drosophila antibody (pAb)	Raised against amino acids 372-385 of Drosophila GAGA.	Dr	WB	100 µl	39024
GATA-1 antibody (pAb)	Raised against amino acids 66-78 of human GATA-1.	H, M, R	WB	100 µl	39025
GATA-1 antibody (pAb)	Raised against amino acids 66-78 of human GATA-1.	H	EMSA	17 rxns	39319
GATA-4 antibody (pAb)	Raised against a peptide to the C-terminal region of human GATA-4.	H	WB	100 µl 10 µl	39893 39894
NEW GATA-6 antibody (pAb)	Raised against a peptide to the N-terminal region of human GATA-6.	H	WB	100 µl 10 µl	61063 61064
GCN5 antibody (pAb)	Raised against a peptide to the N-terminal region of human GCN5.	H	WB	100 µl 10 µl	39831 39832

Reactivity: B = Bovine; BY = Budding Yeast; Ce = *C. elegans*; Dr = *Drosophila*; FY = Fission Yeast; Hm = Hamster; H = Human; Mk = Monkey; M = Mouse; NS = Not Species Specific; Rb = Rabbit; R = Rat; WR = Wide range predicted
Applications: ChIP = Chromatin IP; DB = Dot Blot; FC = Flow Cytometry; ICC = Immunocytochemistry; IF = Immunofluorescence; IHC = Immunohistochemistry; IP = Immunoprecipitation; MeDIP = Methyl-DNA IP; WB = Western Blot

Antibody	Immunogen	Reactivity	Applications	Format	Catalog No.
GCN5 antibody (mAb) (Clone 2GC-2C11)	Raised against a peptide to amino acids 18-132 of human GCN5.	H	ELISA, IF, WB	100 µg 10 µg	39975 39976
HA Tag antibody (mAb) (Clone 3B9)	Raised against synthetic peptide containing multiple repeats of the HA tag epitope, YPYDVPDYA.	WR	IP	200 µg 10 µg	39627 39628
Hbo1 antibody (pAb)	Raised against a peptide to amino acids 158-172 of human Hbo1.	H, WR	WB	200 µl 10 µl	39105 39106
HBP-1 antibody (mAb)	Raised against full-length human HBP-1.	H, R	ChIP, IF, WB	100 µl	39511
HDAC1 antibody (mAb) (Clone 10E2)	Raised against a KLH-peptide to amino acids 467-482 of human and murine HDAC1.	H, M, R	ChIP, IF, IHC, IP, WB	200 µl 10 µl	39531 39532
HDAC1 antibody (mAb) (Clone 10E2)	Raised against a peptide to amino acids 467-482 of murine HDAC1.	H, M, R	WB	100 µg 10 µg	39881 39882
HDAC1 antibody (pAb)	Peptide mixture of amino acids 1-5, 433-448, 467-482 of human HDAC1.	H	WB	100 µg	40967
HDAC2 antibody (mAb) (Clone 3F3)	Raised against a KLH-peptide to amino acids 541-556 of human HDAC2.	H, M, R	ChIP, IF, IHC, IP, WB	200 µl 10 µl	39533 39534
HDAC2 antibody (mAb) (Clone 3F3)	Raised against a peptide to amino acids 541-556 of human HDAC2.	H, M, R	WB	100 µg 10 µg	39883 39884
HDAC3 antibody (pAb)	Raised against amino acids 2-17 of human HDAC3.	H	ChIP, WB	100 µg	40968
HDAC3 antibody (mAb) (Clone 3G6)	Raised against amino acids 411-428 of human HDAC3.	H	IHC, IP, WB	100 µg 10 µg	39717 39718
HDAC4 antibody (pAb)	Raised against amino acids 194-209 of human HDAC4.	H	ChIP, WB	100 µg	40969
HDAC5 antibody (pAb)	Raised against amino acids 572-587 of human HDAC5.	H, M	ChIP, WB	100 µg	40970
HDAC6 antibody (pAb)	Raised against amino acids 1-16 of human HDAC6.	H, M	ChIP, WB	100 µg	40971
HDAC9 antibody (pAb)	Raised against a peptide containing the C-terminal region of human HDAC9.	H	WB	100 µl 10 µl	39793 39794
HDAC11 antibody (pAb)	Raised against amino acids 238-251 of human HDAC11.	H, M	WB	200 µl	39208
HIF-1 alpha antibody (pAb)	Raised against a peptide to the N-terminal region of human HIF-1 alpha.	H	WB	100 µl 10 µl	39665 39666
HIPK1 antibody (mAb)	Raised against a recombinant protein including amino acids 702-925 of mouse HIPK1.	M	IF	100 µg 10 µg	39725 39726
HIPK2 antibody (mAb)	Raised against a recombinant protein including amino acids 898-1051 of mouse HIPK2.	H	IF, IHC, IP, WB	100 µg 10 µg	39677 39678
HIRA antibody (mAb) (Clone WC15)	Raised against amino acids 421-729 of GST-tagged human HIRA.	H, M	IP	200 µg 10 µg	39555 39556
HIRA antibody (mAb) (Clone WC119.2H11)	Raised against amino acids 421-729 of GST-tagged human HIRA.	H, M	WB	200 µg 10 µg	39557 39558
Histone H1 antibody (pAb)	Raised against calf thymus histone H1.	H, WR	IF, WB	100 µl 10 µl	39707 39708

Reactivity: B = Bovine; BY = Budding Yeast; Ce = *C. elegans*; Dr = *Drosophila*; FY = Fission Yeast; Hm = Hamster; H = Human; Mk = Monkey; M = Mouse; NS = Not Species Specific; Rb = Rabbit; R = Rat; WR = Wide range predicted
Applications: ChIP = Chromatin IP; DB = Dot Blot; FC = Flow Cytometry; ICC = Immunocytochemistry; IF = Immunofluorescence; IHC = Immunohistochemistry; IP = Immunoprecipitation; MeDIP = Methyl-DNA IP; WB = Western Blot

Antibody	Immunogen	Reactivity	Applications	Format	Catalog No.
Histone H1 antibody (pAb)	Raised against recombinant <i>Drosophila</i> histone H1.	Dr	IF, IP, WB	100 µl	39575
Histone H2A antibody (pAb)	Raised against recombinant yeast histone H2A.	BY, FY	ChIP, WB	200 µl 10 µl	39235 39236
Histone H2A antibody (pAb)	Raised against recombinant yeast histone H2A.	BY, FY	WB	100 µg 10 µg	39945 39946
Histone H2A antibody (pAb)	Peptide mixture of amino acids 1-5 and 81-96 of human histone H2A.	H	WB	100 µg	39209
Histone H2A, acidic patch antibody (pAb)	Raised against a peptide with amino acids 89-91 of human histone H2A.	H, WR	ChIP, WB	200 µl 10 µl	39111 39112
Histone H2A, C-terminal antibody (pAb)	Raised against a peptide with the C-terminal region of human histone H2A.	H, WR	WB	200 µl 10 µl	39591 39592
Histone H2A acetyl Lys5 antibody (pAb)	Raised against a peptide with acetyl-lysine 5 of human histone H2A.	H, WR	WB	200 µl 10 µl	39107 39108
Histone H2A acetyl Lys9 antibody (pAb)	Raised against a peptide with acetyl-lysine 9 of human histone H2A.	H, WR	WB	200 µl 10 µl	39109 39110
Histone H2A phospho Thr120 antibody (pAb)	Raised against a peptide with phospho-threonine 120 of human histone H2A.	H, WR	WB	200 µl 10 µl	39391 39392
Histone H2A phospho Ser129 antibody (pAb)	Raised against peptide with phospho-serine 129 of budding yeast H2A.	BY	ChIP, IF, IP, WB	200 µl 10 µl	39271 39272
Histone macroH2A1 antibody (pAb)	Raised against the N-terminal half of recombinant mouse macroH2A1.	H, M	IF, WB	200 µl 10 µl	39593 39594
Histone macroH2A1.1 antibody (pAb)	Raised against a peptide to amino acids 198-226 of human macroH2A1.1.	H	IF, IHC, WB	100 µg 10 µg	39871 39872
Histone macroH2A2 antibody (pAb)	Raised against a peptide to amino acids 162-372 of human macroH2A2.	H	IF, IHC, WB	100 µg 10 µg	39873 39874
Histone H2A/H4 phospho Ser1 antibody (pAb)	Raised against a peptide with phospho-serine 1 of human histone H2A.	H, WR	WB	200 µl 10 µl	39115 39116
Histone H2Av antibody (pAb)	Raised against a peptide in the C-terminus of the <i>Drosophila melanogaster</i> histone variant H2Av.	Dr	WB	100 µl 10 µl	39715 39716
Histone H2AX antibody (pAb)	Raised against a peptide to the C-terminus of human histone H2AX.	H	IF, WB	100 µl 10 µl	39689 39690
Histone H2AX phospho Ser139 antibody (pAb)	Raised against a peptide with phospho-serine 139 of human histone H2AX.	H, WR	IF, WB	200 µg 10 µg	39117 39118
Histone H2A.Z antibody (pAb)	Raised against a peptide to the C-terminus of human histone H2A.Z.	H, WR	ChIP, WB	200 µl 10 µl	39113 39114
Histone H2A.Z antibody (pAb)	Raised against a peptide to the C-terminus of human histone H2A.Z.	H, WR	WB	100 µg 10 µg	39943 39944
Htz1 antibody (pAb)	Raised against full-length Htz1 protein, the <i>S. cerevisiae</i> homolog of Histone H2A.Z.	BY	ChIP, WB	100 µl 10 µl	39647 39648

Reactivity: B = Bovine; BY = Budding Yeast; Ce = *C. elegans*; Dr = *Drosophila*; FY = Fission Yeast; Hm = Hamster; H = Human; Mk = Monkey; M = Mouse; NS = Not Species Specific; Rb = Rabbit; R = Rat; WR = Wide range predicted
Applications: ChIP = Chromatin IP; DB = Dot Blot; FC = Flow Cytometry; ICC = Immunocytochemistry; IF = Immunofluorescence; IHC = Immunohistochemistry; IP = Immunoprecipitation; MeDIP = Methyl-DNA IP; WB = Western Blot

Antibody	Immunogen	Reactivity	Applications	Format	Catalog No.
Pht1 / H2AZ antibody (pAb)	Raised against a peptide with the C-terminus of Pht1, the Histone H2A variant in <i>S. pombe</i> .	FY	WB	100 µl 10 µl	39640 39641
Pht1 / H2AZ acetyl antibody (pAb)	Raised against a peptide with acetyl-lysine at positions 5, 7, 12 and 16 from the amino terminus of Pht1, the Histone H2A variant in <i>S. pombe</i> .	FY	WB	100 µl 10 µl	39642 39643
Histone H2B antibody (pAb)	Raised against a peptide with the C-terminus of human histone H2B.	H, WR	WB	200 µl 10 µl	39125 39126
Histone H2B antibody (pAb)	Raised against recombinant yeast histone H2B.	BY, FY	ChIP, WB	200 µl 10 µl	39237 39238
Histone H2B antibody (pAb)	Raised against recombinant yeast histone H2B.	BY, FY	WB	100 µg 10 µg	39947 39948
Histone H2B antibody (pAb)	Raised against amino acids 111-125 of human histone H2B.	H	WB	100 µg	39210
NEW Histone H2B antibody (mAb) (Clone 5HH2-2A8)	Raised against a peptide with the C-terminus of human histone H2B.	H	WB	100 µg 10 µg	61037 61038
Histone H2B acetyl Lys5 antibody (pAb)	Raised against a peptide with acetyl-lysine 5 of human histone H2B.	H, WR	ChIP, WB	200 µl 10 µl	39123 39124
Histone H2B acetyl Lys12 antibody (pAb)	Raised against a peptide with acetyl-lysine 12 of human histone H2B.	H, WR	ChIP, WB	100 µl 10 µl	39669 39670
NEW Histone H2B phospho Ser14 antibody (mAb) (Clone MABI 0251)	Raised against a peptide with phospho-serine 14 of human histone H2B.	H, WR	IF, IP, WB	100 µg	61011
Histone H2B acetyl Lys16 antibody (pAb)	Raised against a peptide with acetyl-lysine 16 of human histone H2B.	H, WR	ChIP, WB	200 µl 10 µl	39121 39122
Histone H2B acetyl Lys46 antibody (pAb)	Raised against a peptide with acetyl-lysine 46 of human histone H2B.	H, WR	ChIP, WB	200 µl 10 µl	39571 39572
Histone H2B dimethyl Lys46 antibody (pAb)	Peptide containing dimethyl-lysine 46 of human histone H2B.	H, WR	IF, WB	200 µl 10 µl	39567 39568
Histone H2B acetyl Lys120 antibody (pAb)	Raised against a peptide with acetyl-lysine 120 of human histone H2B.	H, WR	ChIP, WB	200 µl 10 µl	39119 39120
Histone H2B ubiquityl Lys120 antibody (mAb) (Clone 56)	Raised against a branched peptide corresponding the the ubiquitin conjugation site at Lys120 of human Histone H2B.	H, WR	ChIP, WB	100 µg 10 µg	39623 39624
Histone H3 antibody (mAb) (Clone MABI 0301)	Raised against an N-terminal peptide of human histone H3.	H, WR	ChIP, IF, WB	100 µg	39763
Histone H3, C-terminal antibody (pAb)	Raised against a C-terminal peptide of human histone H3.	BY, H, WR	ChIP, WB	200 µl 10 µl	39163 39164
Histone H3.cs1 antibody (pAb)	Raised against peptide with a Cathepsin L cleavage site at N-terminal histone H3.	H	WB	200 µl 10 µl	39573 39574
Histone H3 acetyl antibody (pAb)	Raised against peptide with acetyl-lysines at the H3 N-terminal tail.	H, WR	ChIP, WB	200 µl 10 µl	39139 39140
Histone H3 dimethyl Arg2 symmetric antibody (pAb)	Raised against a peptide with dimethyl-arginine 2 (symmetric) of human histone H3.	H, WR	WB	100 µl 10 µl	39703 39704

Reactivity: B = Bovine; BY = Budding Yeast; Ce = *C. elegans*; Dr = *Drosophila*; FY = Fission Yeast; Hm = Hamster; H = Human; Mk = Monkey; M = Mouse; NS = Not Species Specific; Rb = Rabbit; R = Rat; WR = Wide range predicted
Applications: ChIP = Chromatin IP; DB = Dot Blot; FC = Flow Cytometry; ICC = Immunocytochemistry; IF = Immunofluorescence; IHC = Immunohistochemistry; IP = Immunoprecipitation; MeDIP = Methyl-DNA IP; WB = Western Blot

Antibody	Immunogen	Reactivity	Applications	Format	Catalog No.
Histone H3 phospho Thr3 antibody (pAb)	Raised against a peptide with phospho-threonine 3 of human histone H3.	H, WR	WB	200 µl	39153
				10 µl	39154
Histone H3 acetyl Lys4 antibody (pAb)	Raised against a peptide with acetyl-lysine 4 of human histone H3.	H, WR	ChIP, IF, WB	200 µl	39381
				10 µl	39382
Histone H3 monomethyl Lys4 antibody (mAb) (Clone MABI 0302)	Raised against a peptide with monomethyl-lysine 4 of human histone H3.	H, WR	ChIP, WB	100 µg	39635
Histone H3 monomethyl Lys4 antibody (pAb)	Raised against peptide with monomethyl-lysine 4 of human histone H3.	H, WR	IF, WB	200 µl	39297
				10 µl	39298
Histone H3 dimethyl Lys4 antibody (mAb) (Clone MABI 0303)	Raised against a peptide with dimethyl-lysine 4 of human histone H3.	H, WR	ChIP, IF, WB	100 µg	39679
Histone H3 dimethyl Lys4 antibody (pAb)	Raised against a peptide with dimethyl-lysine 4 of human histone H3.	H, WR	ChIP, WB	200 µl	39141
				10 µl	39142
Histone H3 dimethyl Lys4 antibody (pAb)	Raised against a peptide with dimethyl-lysine 4 of human histone H3.	H, WR	WB	100 µg	39913
				10 µg	39914
Histone H3 trimethyl Lys4 antibody (pAb)	Raised against a peptide with trimethyl-lysine 4 of human histone H3.	BY, H, WR	ChIP, WB	200 µl	39159
				10 µl	39160
Histone H3 trimethyl Lys4 antibody (pAb)	Raised against a peptide with trimethyl-lysine 4 of human histone H3.	BY, H, WR	WB	100 µg	39915
				10 µg	39916
Histone H3 monomethyl Arg8 antibody (pAb)	Raised against a peptide with monomethyl-arginine 8 of human histone H3.	H, WR	WB	100 µl	39673
				10 µl	39674
Histone H3 dimethyl Arg8 asymmetric antibody (pAb)	Raised against a peptide with dimethyl-arginine 8 (asymmetric) of human histone H3.	H	WB	100 µl	39651
				10 µl	39652
Histone H3 acetyl Lys9 antibody (pAb)	Raised against a peptide with acetyl-lysine 9 of human histone H3.	H, WR	ChIP, WB	200 µl	39137
				10 µl	39138
Histone H3 acetyl Lys9 antibody (pAb)	Raised against a peptide with acetyl-lysine 9 of human histone H3.	H, WR	IF, WB	200 µl	39585
				10 µl	39586
Histone H3 acetyl Lys9 antibody (pAb)	Raised against a peptide with acetyl-lysine 9 of human histone H3.	H, WR	WB	100 µg	39917
				10 µg	39918
Histone H3 pan-methyl Lys9 antibody (pAb)	Raised against a peptide with methyl-lysine 9 of human histone H3.	H, WR	IF, WB	200 µl	39241
				10 µl	39242
Histone H3 monomethyl Lys9 antibody (mAb) (Clone MABI 0306)	Raised against a peptide with monomethyl-lysine 9 of human histone H3.	H, WR	ChIP, WB	100 µg	39681
Histone H3 monomethyl Lys9 antibody (pAb)	Raised against a peptide with monomethyl-lysine 9 of human histone H3.	H, WR	IF, WB	200 µl	39249
				10 µl	39250
Histone H3 monomethyl Lys9 antibody (pAb)	Raised against a peptide with monomethyl-lysine 9 of human histone H3.	H, WR	WB	100 µg	39887
				10 µg	39888
Histone H3 dimethyl Lys9 antibody (mAb) (Clone MABI 0307)	Raised against a peptide with dimethyl-lysine 9 of human histone H3.	H, WR	ChIP, IF, WB	100 µg	39683

Reactivity: B = Bovine; BY = Budding Yeast; Ce = *C. elegans*; Dr = *Drosophila*; FY = Fission Yeast; Hm = Hamster; H = Human; Mk = Monkey; M = Mouse; NS = Not Species Specific; Rb = Rabbit; R = Rat; WR = Wide range predicted
Applications: ChIP = Chromatin IP; DB = Dot Blot; FC = Flow Cytometry; ICC = Immunocytochemistry; IF = Immunofluorescence; IHC = Immunohistochemistry; IP = Immunoprecipitation; MeDIP = Methyl-DNA IP; WB = Western Blot

Antibody	Immunogen	Reactivity	Applications	Format	Catalog No.
Histone H3 dimethyl Lys9 antibody (pAb)	Raised against a peptide with dimethyl-lysine 9 of human histone H3.	H, M, WR	ChIP, IF, WB	200 µl	39239
				10 µl	39240
Histone H3 dimethyl Lys9 antibody (pAb)	Raised against a peptide with dimethyl-lysine 9 of human histone H3.	H, WR	ChIP, IF, WB	200 µl	39375
				10 µl	39376
Histone H3 dimethyl Lys9 antibody (pAb)	Raised against a peptide with dimethyl-lysine 9 of human histone H3.	H, WR	ChIP, IF, WB	100 µg	39753
				10 µg	39754
NEW Histone H3 trimethyl Lys9 antibody (mAb) (Clone MABI 0319)	Raised against peptide with trimethyl-lysine 9 of human histone H3.	H, WR	ChIP, IF, IP, WB	100 µg	61013
Histone H3 trimethyl Lys9 antibody (pAb)	Raised against a peptide with trimethyl-lysine 9 of human histone H3.	FY, H, WR	ChIP, WB	200 µl	39161
				10 µl	39162
Histone H3 trimethyl Lys9 antibody (mAb) (Clone 2AG-6F12-H4)	Raised against a peptide with trimethyl-lysine 9 of human histone H3.	H, WR	IF, IP, WB	200 µl	39285
				10 µl	39286
Histone H3 trimethyl Lys9 antibody (pAb)	Raised against a peptide with trimethyl-lysine 9 of human histone H3.	H, WR	IF, WB	100 µg	39765
				10 µg	39766
Histone H3 phospho Ser10 antibody (mAb) (Clone MABI 0312)	Raised against a peptide with phospho-serine 10 of human histone H3.	H, WR	IF	100 µg	39636
Histone H3 phospho Ser10 antibody (pAb)	Raised against a peptide with phospho-serine 10 of human histone H3.	H, WR	ChIP, IF, WB	200 µl	39253
				10 µl	39254
Histone H3 phospho Ser10,28 antibody (pAb)	Raised against a peptide with phospho-serine 28 of human histone H3.	H, WR	ChIP, IF, WB	200 µl	39147
				10 µl	39148
Histone H3 phospho Thr11 antibody (mAb) (Clone NI23/48)	Raised against a peptide with phospho-threonine 11 of human histone H3.	H	ICC, WB	100 µg	39821
				10 µg	39822
Histone H3 phospho Thr11 antibody (pAb)	Raised against a peptide with phospho-threonine 11 of human histone H3.	H, WR	WB	200 µl	39151
				10 µl	39152
Histone H3 acetyl Lys14 antibody (pAb)	Raised against a peptide with acetyl-lysine 14 of human histone H3.	H, WR	ChIP, WB	200 µl	39599
				10 µl	39616
Histone H3 acetyl Lys14 antibody (pAb)	Raised against a peptide with acetyl-lysine 14 of human histone H3.	H, WR	ChIP, WB	100 µg	39697
				10 µg	39698
Histone H3 dimethyl Lys14 antibody (pAb)	Raised against a peptide with dimethyl-lysine 14 of human histone H3.	H, WR	WB	200 µl	39349
				10 µl	39350
Histone H3 dimethyl Arg17 asymmetric antibody (pAb)	Raised against a peptide with dimethyl-arginine 17 (asymmetric) of human histone H3.	H, WR	WB	100 µg	39709
				10 µg	39710
Histone H3 acetyl Lys18 antibody (pAb)	Raised against a peptide with acetyl-lysine 18 of human histone H3.	H, WR	ChIP, IF, WB	100 µl	39693
				10 µl	39694
Histone H3 acetyl Lys18 antibody (pAb)	Raised against a peptide with acetyl-lysine 18 of human histone H3.	H, WR	ChIP, IF, WB	200 µl	39587
				10 µl	39588
Histone H3 acetyl Lys18 antibody (pAb)	Raised against a peptide with acetyl-lysine 18 of human histone H3.	H, WR	IF, WB	200 µl	39129
				10 µl	39130

Reactivity: B = Bovine; BY = Budding Yeast; Ce = *C. elegans*; Dr = *Drosophila*; FY = Fission Yeast; Hm = Hamster; H = Human; Mk = Monkey; M = Mouse; NS = Not Species Specific; Rb = Rabbit; R = Rat; WR = Wide range predicted
Applications: ChIP = Chromatin IP; DB = Dot Blot; FC = Flow Cytometry; ICC = Immunocytochemistry; IF = Immunofluorescence; IHC = Immunohistochemistry; IP = Immunoprecipitation; MeDIP = Methyl-DNA IP; WB = Western Blot

Antibody	Immunogen	Reactivity	Applications	Format	Catalog No.
Histone H3 acetyl Lys18 antibody (pAb)	Raised against a peptide with acetyl-lysine 18 of human histone H3.	H, WR	ChIP, IF, WB	100 µg 10 µg	39755 39756
Histone H3 monomethyl Lys18 antibody (pAb)	Raised against a peptide with monomethyl-lysine 18 of human histone H3.	H	WB	100 µl 10 µl	39667 39668
Histone H3 acetyl Lys23 antibody (pAb)	Raised against a peptide with acetyl-lysine 23 of human histone H3.	H, WR	ChIP, IF, WB	200 µl 10 µl	39131 39132
Histone H3 monomethyl Lys23 antibody (pAb)	Raised against a peptide with monomethyl-lysine 23 of human histone H3.	H, WR	WB	200 µl 10 µl	39387 39388
Histone H3 dimethyl Lys23 antibody (pAb)	Raised against a peptide with dimethyl-lysine 23 of human histone H3.	H	WB	100 µl 10 µl	39653 39654
Histone H3 acetyl Lys27 antibody (mAb) (Clone MABI 0309)	Raised against a peptide with acetyl-lysine 27 of human histone H3.	H, WR	ChIP, IF, WB	100 µg	39685
Histone H3 acetyl Lys27 antibody (pAb)	Raised against a peptide with acetyl-lysine 27 of human histone H3.	BY, H, WR	ChIP, IF, WB	200 µg 10 µg	39133 39134
Histone H3 acetyl Lys27 antibody (pAb)	Raised against a peptide with acetyl-lysine 27 of human histone H3.	BY, H, WR	ChIP, WB	200 µl 10 µl	39135 39136
NEW Histone H3 monomethyl Lys27 antibody (mAb) (Clone MABI 0321)	Raised against peptide with monomethyl-lysine 27 of human histone H3.	H, WR	ChIP, IF, IP, WB	100 µg	61015
Histone H3 monomethyl Lys27 antibody (pAb)	Raised against peptide with monomethyl-lysine 27 of human histone H3.	H, WR	IF, WB	200 µl 10 µl	39377 39378
Histone H3 monomethyl Lys27 antibody (pAb)	Raised against peptide with monomethyl-lysine 27 of human histone H3.	H, WR	WB	100 µg 10 µg	39889 39890
Histone H3 dimethyl Lys27 antibody (pAb)	Raised against a peptide with dimethyl-lysine 27 of human histone H3.	H, WR	ChIP, IF, WB	200 µl 10 µl	39245 39246
Histone H3 dimethyl Lys27 antibody (pAb)	Raised against a peptide with dimethyl-lysine 27 of human histone H3.	H, WR	WB	100 µg 10 µg	39919 39920
NEW Histone H3 trimethyl Lys27 antibody (mAb) (Clone MABI 0323)	Raised against peptide with trimethyl-lysine 27 of human histone H3.	H, WR	ChIP, IF, IP, WB	100 µg	61017
Histone H3 di/trimethyl Lys27 antibody (mAb)	Raised against a peptide with trimethyl-lysine 27 of human histone H3.	H, WR	ChIP, WB	200 µl 10 µl	39535 39537
Histone H3 di/trimethyl Lys27 antibody (mAb)	Raised against a peptide with trimethyl-lysine 27 of human histone H3.	H, WR	IF, WB	200 µl 10 µl	39536 39538
Histone H3 trimethyl Lys27 antibody (pAb)	Raised against a peptide with trimethyl-lysine 27 of human histone H3.	H, WR	ChIP, IF, WB	200 µg 10 µg	39155 39157
Histone H3 trimethyl Lys27 antibody (pAb)	Raised against a peptide with trimethyl-lysine 27 of human histone H3.	H, WR	ChIP, WB	200 µl 10 µl	39156 39158
Histone H3 phospho Ser28 antibody (mAb) (Clone HTA28)	Raised against a peptide with phospho-serine 28 of human histone H3.	H, M, R	IF, WB	100 µl	39098

Reactivity: B = Bovine; BY = Budding Yeast; Ce = *C. elegans*; Dr = *Drosophila*; FY = Fission Yeast; Hm = Hamster; H = Human; Mk = Monkey; M = Mouse; NS = Not Species Specific; Rb = Rabbit; R = Rat; WR = Wide range predicted
Applications: ChIP = Chromatin IP; DB = Dot Blot; FC = Flow Cytometry; ICC = Immunocytochemistry; IF = Immunofluorescence; IHC = Immunohistochemistry; IP = Immunoprecipitation; MeDIP = Methyl-DNA IP; WB = Western Blot

Antibody	Immunogen	Reactivity	Applications	Format	Catalog No.
Histone H3 phospho Ser28 antibody (pAb)	Raised against a peptide with phospho-serine 28 of human histone H3.	H, WR	WB	200 µl	39149
				10 µl	39150
Histone H3.3 phospho Ser31 antibody (pAb)	Raised against a peptide with phospho-serine 31 of human histone H3.3.	H	WB	200 µl	39637
				10 µl	39638
Histone H3 acetyl Lys36 antibody (pAb)	Raised against a peptide with acetyl-lysine 36 of human histone H3.	H, WR	ChIP, IF, WB	200 µl	39379
				10 µl	39380
NEW Histone H3 dimethyl Lys36 antibody (mAb) (Clone MABI 0332)	Raised against a peptide with dimethyl-lysine 36 of human histone H3.	H, WR	ChIP, IF, IP, WB	100 µg	61019
Histone H3 dimethyl Lys36 antibody (pAb)	Raised against a peptide with dimethyl-lysine 36 of human histone H3.	BY, H, WR	ChIP, IF, WB	200 µl	39255
				10 µl	39256
Histone H3 dimethyl Lys36 antibody (pAb)	Raised against a peptide with dimethyl-lysine 36 of human histone H3.	BY, H, WR	WB	100 µg	39891
				10 µg	39892
NEW Histone H3 trimethyl Lys36 antibody (mAb) (Clone MABI 0333)	Raised against peptide with trimethyl-lysine 36 of human histone H3.	H, WR	ChIP, IF, IP, WB	100 µg	61021
Histone H3 phospho Thr45 antibody (pAb)	Raised against a peptide with phospho-threonine 45 of human histone H3.	BY, H, WR	WB	100 µl	39737
				10 µl	39738
NEW Histone H3 acetyl Lys56 antibody (mAb) (Clone 12.1)	Raised against peptide with acetyl-lysine 56 of yeast histone H3.	BY, FY, H	ChIP, DB	100 µg	61061
				10 µg	61062
Histone H3 acetyl Lys56 antibody (pAb)	Raised against peptide with acetyl-lysine 56 of yeast histone H3.	BY, H	ChIP, WB	200 µl	39281
				10 µl	39282
Histone H3 monomethyl Lys56 antibody (pAb)	Raised against peptide with monomethyl-lysine 56 of human histone H3.	H	WB	200 µl	39273
				10 µl	39274
Histone H3 dimethyl Lys56 antibody (pAb)	Raised against a peptide with dimethyl-lysine 56 of human histone H3.	H	WB	200 µl	39277
				10 µl	39278
Histone H3 acetyl Lys64 antibody (pAb)	Raised against a peptide with acetyl-lysine 64 of human histone H3.	BY, H, WR	ChIP, IF, WB	200 µl	39545
				10 µl	39546
Histone H3 acetyl Lys79 antibody (pAb)	Peptide containing acetyl-lysine 79 of human histone H3.	H, WR	ChIP, WB	200 µl	39565
				10 µl	39566
Histone H3 monomethyl Lys79 antibody (pAb)	Raised against a peptide with monomethyl-lysine 79 of human histone H3.	H, WR	ChIP, WB	200 µl	39145
				10 µl	39146
Histone H3 monomethyl Lys79 antibody (pAb)	Raised against a peptide with monomethyl-lysine 79 of human histone H3.	H, WR	WB	100 µg	39921
				10 µg	39922
Histone H3 dimethyl Lys79 antibody (pAb)	Raised against a peptide with dimethyl-lysine 79 of human histone H3.	H, WR	ChIP, WB	200 µl	39143
				10 µl	39144
Histone H3 dimethyl Lys79 antibody (pAb)	Raised against a peptide with dimethyl-lysine 79 of human histone H3.	H, WR	WB	100 µg	39923
				10 µg	39924
Histone H3 monomethyl Lys122 antibody (pAb)	Raised against a peptide with monomethyl-lysine 122 of human H3.	H, WR	WB	200 µl	39367
				10 µl	39368

Reactivity: B = Bovine; BY = Budding Yeast; Ce = *C. elegans*; Dr = *Drosophila*; FY = Fission Yeast; Hm = Hamster; H = Human; Mk = Monkey; M = Mouse; NS = Not Species Specific; Rb = Rabbit; R = Rat; WR = Wide range predicted
Applications: ChIP = Chromatin IP; DB = Dot Blot; FC = Flow Cytometry; ICC = Immunocytochemistry; IF = Immunofluorescence; IHC = Immunohistochemistry; IP = Immunoprecipitation; MeDIP = Methyl-DNA IP; WB = Western Blot

Antibody	Immunogen	Reactivity	Applications	Format	Catalog No.
Histone H4 antibody (pAb)	Raised against recombinant human histone H4.	H, WR	WB	200 µl	39269
				10 µl	39270
Histone H4 pan-acetyl antibody (pAb)	Raised against a peptide of Tetrahymena histone H2A acetylated at multiple lysines.	H, WR	ChIP, IF, WB	200 µl	39243
				10 µl	39244
Histone H4 pan-acetyl antibody (pAb)	Raised against a peptide of Tetrahymena histone H2A acetylated at multiple lysines.	H, WR	WB	100 µg	39925
				10 µg	39926
Histone H4 tetra-acetyl antibody (pAb)	Raised against a peptide of Tetrahymena hvl acetylated at multiple lysines.	BY, H, WR	IF, WB	50 µg	39177
				10 µg	39227
Histone H4 tetra-acetyl antibody (pAb)	Raised against a peptide of Tetrahymena hvl acetylated at multiple lysines.	BY, H, WR	ChIP, WB	50 µl	39179
				10 µl	39228
Histone H4 acetyl antibody (mAb) (Clone 3HH4-2C2)	Raised against a peptide to amino acids 1- 24 of human histone H4 acetylated at lysines 5, 8, 12 and 16.	H, WR	ELISA, ICC, IP, WB	100 µg	39967
				10 µg	39968
Histone H4 dimethyl Arg3, asymmetric antibody (pAb)	Raised against a peptide with dimethyl-arginine 3 (asymmetric) of human histone H4.	H, WR	IF, WB	100 µl	39705
				10 µl	39706
Histone H4 acetyl Lys5 antibody (pAb)	Raised against a peptide with acetyl-lysine 5 of human histone H4.	H, WR	ChIP, IF, WB	200 µl	39169
				10 µl	39170
Histone H4 acetyl Lys5 antibody (pAb)	Raised against a peptide with acetyl-lysine 5 of human histone H4.	H, WR	ChIP, IF, WB	200 µl	39583
				10 µl	39584
Histone H4 acetyl Lys5 antibody (pAb)	Raised against a peptide with acetyl-lysine 5 of human histone H4.	H, WR	ChIP, WB	100 µg	39699
				10 µg	39700
Histone H4 acetyl Lys8 antibody (pAb)	Raised against a peptide with acetyl-lysine 8 of human histone H4.	H, WR	WB	200 µl	39171
				10 µl	39172
Histone H4 acetyl Lys12 antibody (pAb)	Raised against a peptide with acetyl-lysine 12 of human histone H4.	BY, H, WR	ChIP, IF, WB	200 µl	39165
				10 µl	39166
Histone H4 acetyl Lys12 antibody (pAb)	Raised against a peptide with acetyl-lysine 12 of human histone H4.	BY, H, WR	WB	100 µg	39927
				10 µg	39928
Histone H4 acetyl Lys16 antibody (pAb)	Raised against a peptide with acetyl-lysine 16 of human histone H4.	BY, Dr, H, WR	ChIP, WB	200 µl	39167
				10 µl	39168
Histone H4 acetyl Lys16 antibody (pAb)	Raised against a peptide with acetyl-lysine 16 of human histone H4.	BY, Dr, H, WR	WB	100 µg	39929
				10 µg	39930
Histone H4 monomethyl Lys20 antibody (mAb) (Clone 5E10-D8)	Raised against a peptide with monomethyl-lysine 20 of human histone H4.	H, WR	IF, WB	100 µg	39727
				10 µg	39728
Histone H4 monomethyl Lys20 antibody (pAb)	Raised against a peptide with monomethyl-lysine 20 of human histone H4.	H, WR	ChIP, IF, WB	200 µl	39175
				10 µl	39176
Histone H4 dimethyl Lys20 antibody (mAb)	Raised against a peptide with dimethyl-lysine 20 of human histone H4.	H, WR	WB	200 µl	39539
				10 µl	39540
Histone H4 dimethyl Lys20 antibody (pAb)	Raised against a peptide with dimethyl-lysine 20 of human histone H4.	H, WR	ChIP, IF, WB	200 µl	39173
				10 µl	39174

Reactivity: B = Bovine; BY = Budding Yeast; Ce = *C. elegans*; Dr = *Drosophila*; FY = Fission Yeast; Hm = Hamster; H = Human; Mk = Monkey; M = Mouse; NS = Not Species Specific; Rb = Rabbit; R = Rat; WR = Wide range predicted
Applications: ChIP = Chromatin IP; DB = Dot Blot; FC = Flow Cytometry; ICC = Immunocytochemistry; IF = Immunofluorescence; IHC = Immunohistochemistry; IP = Immunoprecipitation; MeDIP = Methyl-DNA IP; WB = Western Blot

Antibody	Immunogen	Reactivity	Applications	Format	Catalog No.
Histone H4 trimethyl Lys20 antibody (mAb) (Clone 6F8-D9)	Raised against a peptide with trimethyl-lysine 20 of human histone H4.	H, M, WR	ChIP, WB	100 µg	39671
				10 µg	39672
Histone H4 trimethyl Lys20 antibody (pAb)	Raised against a peptide with trimethyl-lysine 20 of human histone H4.	H, WR	ChIP, IF, WB	200 µl	39180
				10 µl	39181
Histone H4 monomethyl Lys31 antibody (pAb)	Raised against a peptide with monomethyl lysine 31 of human histone H4.	H, WR	WB	200 µl	39385
				10 µl	39386
HMG-1 antibody (pAb)	Raised against amino acids 167-182 of human HMG-1.	H	WB	200 µl	39551
				10 µl	39552
HMG-2 antibody (pAb)	Raised against amino acids 166-181 of human HMG-2.	H	WB	100 µl	39029
HMGA1 antibody (pAb)	Raised against a peptide containing the N-terminal region of human HMGA1.	H	IF, WB	100 µl	39615
				10 µl	39650
NEW HMGA2 / HMG1-C antibody (pAb)	Raised against a peptide containing the N-terminal region of human HMGA2	H	WB	100 µl	61041
				10 µl	61042
HMGNI antibody (pAb)	Raised against a peptide containing the C-terminal region of human HMGNI.	H	WB	100 µl	39833
				10 µl	39834
HNF-1α antibody (pAb)	Raised against amino acids 540-555 of human HNF-1α.	H, M, R	WB	100 µl	39030
HNF-1α antibody (pAb)	Raised against amino acids 540-555 of human HNF-1α.	H, M, R	EMSA	17 rxns	39321
HNF-3α / FOXA1 antibody (mAb)	Raised against a recombinant protein with amino acids 7-103 of rat HNF-3α / FOXA1.	H, M	IHC, WB	100 µg	39837
				10 µg	39838
HNF-3β antibody (pAb)	Raised against amino acids 295-311 of human HNF-3β.	H	WB	100 µl	39096
HNF-3β / FOXA2 antibody (pAb)	Raised against a recombinant protein with amino acids 7-86 of rat HNF-3β / FOXA2.	H	IHC, WB	100 µl	39827
				10 µl	39828
HOXA9 antibody (pAb)	Raised against a peptide containing the N-terminal region of human HOXA9.	H	WB	100 µl	39825
				10 µl	39826
HP1α antibody (pAb)	Raised against a recombinant protein to full-length mouse HP1α.	H, M	WB	200 µl	39295
				10 µl	39296
HP1α antibody (mAb) (Clone 2HP-1H5)	Raised against a recombinant protein with amino acids 67-119 of mouse HP1α.	H, M	ChIP, ELISA, ICC, IF, ICH	100 µg	39977
				10 µg	39978
HP1β antibody (mAb) (Clone 1MOD-1A9)	Raised against a recombinant mouse HP1β.	H, M	ELISA, IF, ICH, IP, WB	100 µg	39979
				10 µg	39980
HP1γ antibody (mAb) (Clone 2MOD-1G6)	Raised against a recombinant full-length mouse HP1γ.	H, M	ELISA, ICC, IF, ICH, IP, WB	100 µg	39981
				10 µg	39982
HP1 gamma phospho Ser93 antibody (pAb)	Raised against a peptide with phospho-serine 93 of human HP1 gamma.	H	WB	100 µl	39655
				10 µl	39656
Ikaros antibody (mAb)	Raised against the N-terminal half of mouse Ikaros.	M	WB	200 µl	39355
				10 µl	39356
Ikaros antibody (pAb)	Raised against the N-terminal half of mouse Ikaros.	M	WB	200 µl	39291
				10 µl	39292

Reactivity: B = Bovine; BY = Budding Yeast; Ce = *C. elegans*; Dr = *Drosophila*; FY = Fission Yeast; Hm = Hamster; H = Human; Mk = Monkey; M = Mouse; NS = Not Species Specific; Rb = Rabbit; R = Rat; WR = Wide range predicted
Applications: ChIP = Chromatin IP; DB = Dot Blot; FC = Flow Cytometry; ICC = Immunocytochemistry; IF = Immunofluorescence; IHC = Immunohistochemistry; IP = Immunoprecipitation; MeDIP = Methyl-DNA IP; WB = Western Blot

Antibody	Immunogen	Reactivity	Applications	Format	Catalog No.
IκBα antibody (mAb)	Raised against amino acids 32-391 of recombinant human IκBα.	H, M	IP, WB	100 µg	40903
IκBα phospho Ser32,36 antibody (mAb)	Raised against a peptide with phospho-serine 32 and 36 of human IκBα.	H	WB	100 µg	40904
IKKα antibody (mAb)	Raised against full-length human IKKα.	H, M	IP, WB	100 µg	40905
IKKβ antibody (mAb) (Clone 10AG2)	Raised against 6xHis-tagged, full-length human IKKβ.	H	WB	100 µg	40906
IKKβ antibody (mAb) (Clone 10A9B6)	Raised against 6xHis-tagged, full-length human IKKβ.	H	IP, WB	100 µg	40907
IKKγ antibody (mAb)	Raised against full-length human IKKγ.	H	WB	100 µg	40908
IKKi/IKKε antibody (mAb)	Raised against amino acids 175-188 of human IKKi/IKKε.	H, M, R	WB	100 µg	40950
IKKi/IKKε antibody (pAb)	Raised against amino acids 175-188, 525-540 & 567-580 of human IKKi/IKKε.	H	WB	100 µg	40957
INCENP antibody (mAb)	Raised against amino acids 231-398 of recombinant human INCENP.	H, M, WR	IF	200 µg 10 µg	39259 39260
INHAT-1/TAF-1α/TAF-1β antibody (pAb)	Peptide mix of amino acids 66-8 & 135-151 of human INHAT-1/TAF-1α/TAF-1β.	H, M, R	WB	200 µl	39202
INHAT-2/pp32 antibody (pAb)	Peptide mix of amino acids 87-103 and 231-246 of human INHAT-2/pp32.	H	WB	200 µl	39213
IRAK-1 antibody (pAb)	Raised against amino acids 700-712 of human IRAK-1.	H	WB	100 µg	40910
IRAK-2 antibody (pAb)	Raised against amino acids 546-564 of human IRAK-2.	H	WB	100 µg	40912
IRF-1 antibody (pAb)	Raised against amino acids 126-140 of human IRF-1.	H	EMSA	17 rxns	39325
IRF-3 antibody (pAb)	Raised against amino acids 306-312 of human IRF-3.	H	WB	200 µl 10 µl	39371 39372
IRF-3 antibody (pAb)	Raised against full-length human IRF-3.	B, H	ChIP, WB	100 µl	39033
IRF-6 antibody (pAb)	Raised against amino acids 358-372 of human IRF-6.	H	WB	100 µg	39080
IRF-7 antibody (pAb)	Raised against amino acids 463-477 of human IRF-7.	H	WB	100 µg	39081
JARID1C antibody (pAb)	Raised against a peptide to the C-terminus of human JARID1C.	H, M	WB	200 µl 10 µl	39229 39230
Jhd2 antibody (pAb)	Raised against a peptide to the C-terminus of <i>S. cerevisiae</i> Jhd2.	BY	WB	100 µl 10 µl	39263 39264
JMJD2A antibody (mAb) (Clone NI54/32)	Raised against a recombinant protein containing amino acids 586-674 of human JMJD2A.	H, M	WB	100 µg 10 µg	39815 39816
JMJD2D antibody (pAb)	Raised against amino acids 337-359 of human JMJD2D.	H	WB	200 µl 10 µl	39247 39248
JMJD2F antibody (pAb)	Raised against a peptide to the C-terminus of human JMJD2F.	H	WB	200 µl 10 µl	39257 39258
JunB antibody (pAb)	Raised against amino acids 28-42 of human JunB.	H	WB	200 µl 10 µl	39549 39550
JunB antibody (pAb)	Raised against amino acids 3-17 of human JunB.	H, M	WB	100 µl	39034
JunD antibody (pAb)	Raised against amino acids 143-157 of human JunD.	H	ChIP, EMSA	100 µl	39328
Kaiso antibody (mAb) (Clone 6FID)	Raised against amino acids 103-499 of His-tagged mouse Kaiso.	M	WB	200 µg 10 µg	39365 39366

Reactivity: B = Bovine; BY = Budding Yeast; Ce = *C. elegans*; Dr = *Drosophila*; FY = Fission Yeast; Hm = Hamster; H = Human; Mk = Monkey; M = Mouse; NS = Not Species Specific; Rb = Rabbit; R = Rat; WR = Wide range predicted
Applications: ChIP = Chromatin IP; DB = Dot Blot; FC = Flow Cytometry; ICC = Immunocytochemistry; IF = Immunofluorescence; IHC = Immunohistochemistry; IP = Immunoprecipitation; MeDIP = Methyl-DNA IP; WB = Western Blot

Antibody	Immunogen	Reactivity	Applications	Format	Catalog No.
Ki-67 antibody (pAb)	Raised against a peptide containing the N-terminal region of human Ki-67.	H	WB	100 µl 10 µl	39799 39800
KLF4 antibody (pAb)	Raised against recombinant full-length human KLF4.	H, WR	WB	100 µl 10 µl	39745 39746
L3MBTL1 antibody (pAb)	Raised against a His-tagged fusion protein containing amino acids 1-215 of human L3MBTL2.	H, WR	ChIP, IP, WB	200 µl 10 µl	39182 39183
L3MBTL2 antibody (pAb)	Raised against a His-tagged fusion protein containing amino acids 1-230 of human L3MBTL2.	H	ChIP, IP, WB	200 µl 10 µl	39569 39570
Lamin A, mature antibody (mAb) (Clone 4A4)	Raised against a peptide within the C-terminal region of human Lamin A.	H	IF, IP, WB	100 µg 10 µg	39961 39962
Lamin A/C antibody (mAb) (Clone 3A6-4C11)	Raised against amino acids 430-545 of recombinant human Lamin A.	Hm, H, Mk, M, R	IF, IP, WB	200 µg 10 µg	39287 39288
NEW Lamin A/C R453W antibody (mAb) (Clone 12A-2F5)	Raised against a peptide with amino acids 449-457 of human Lamin A/C and containing the R453W substitution.	H	IF, IP, WB	100 µg 10 µg	39959 39960
Lamin A/C R482W antibody (mAb) (Clone 5H8-B4)	Raised against recombinant protein containing amino acids 430-545 of human Lamin A/C containing the R482W substitution.	H	IF, IP, WB	100 µg 10 µg	39963 39964
Lamin B1 antibody (pAb)	Raised against full-length human Lamin B1.	H, M	WB	100 µl	39095
LAP2 alpha antibody (mAb)	Raised against a peptide with the C-terminus of human LAP2 alpha.	H	WB	200 µl 10 µl	39267 39268
LexA DNA-binding Domain antibody (pAb)	Raised against amino acids 1-222 of recombinant <i>E. coli</i> LexA.	NS	IP, WB	200 µl 10 µl	39184 39185
LIG1 antibody (pAb)	Raised against amino acids 51-68 of human LIG1.	H	WB	200 µl	39214
Livin/ML-IAP antibody (mAb)	Raised against GST-tagged human Livin/ML-IAP.	H	WB	100 µg	40958
LSD1 antibody (pAb)	Raised against full-length recombinant human LSD1 protein.	H, WR	IP, WB	200 µl 10 µl	39186 39187
MAD1 antibody (mAb) (Clone 9B10)	Raised against amino acids 331-718 of mouse MAD1.	H, M	IF, WB	100 µg 10 µg	39659 39660
MAD2L1 antibody (mAb) (Clone 17D10)	Raised against full-length recombinant human MAD2L1.	H, M, WR	WB	200 µg 10 µg	39399 39400
MALT1 antibody (mAb)	Raised against amino acids 701-808 of mouse MALT1 recombinant protein.	H	WB	200 µg 10 µg	39395 39396
Maz antibody (mAb) (Clone 133)	Raised against full-length recombinant human Maz.	H	ChIP, EMSA, IF, IHC, IP, WB	100 µg 10 µg	39935 39936
MBD1 antibody (mAb)	Raised against a peptide corresponding to human MBD1.	H	WB	100 µg	39215
MDB1 antibody (pAb)	Raised against a peptide within the C-terminal region of human MBD1.	H	WB	100 µl 10 µl	39857 39858
MBD2 antibody (pAb)	Raised against a peptide corresponding to human MBD2.	H	WB	200 µl 10 µl	39547 39548

Reactivity: B = Bovine; BY = Budding Yeast; Ce = *C. elegans*; Dr = *Drosophila*; FY = Fission Yeast; Hm = Hamster; H = Human; Mk = Monkey; M = Mouse; NS = Not Species Specific; Rb = Rabbit; R = Rat; WR = Wide range predicted
Applications: ChIP = Chromatin IP; DB = Dot Blot; FC = Flow Cytometry; ICC = Immunocytochemistry; IF = Immunofluorescence; IHC = Immunohistochemistry; IP = Immunoprecipitation; MeDIP = Methyl-DNA IP; WB = Western Blot

Antibody	Immunogen	Reactivity	Applications	Format	Catalog No.
MBD3 antibody (mAb)	Raised against a peptide corresponding to human MBD3.	H	WB	100 µg	39216
MBD4 antibody (pAb)	Peptide mixture of amino acids 268-282 and 337-352 of human MBD4.	H	WB	100 µg	39217
MeCP2 antibody (pAb)	Raised against the N-terminal 320 amino acids of recombinant mouse MeCP2.	H, M	WB	200 µl 10 µl	39188 39189
MeCP2 antibody (pAb)	Peptide mixture of amino acids 11-25 and 181-195 of human MeCP2.	H	WB	100 µg	39218
MeCP2 phospho Ser80 antibody (pAb)	Raised against a peptide with phospho-serine 80 of mouse MeCP2.	H, R	IHC, WB	100 µl 10 µl	39733 39734
NEW MED1 antibody (pAb)	Raised against a peptide to the C-terminal region of human MED1.	H	WB	100 µl 10 µl	61065 61066
MEIS 1/2/3 antibody (mAb) (Clone 9.2.7)	Raised against a peptide containing amino acids 50-390 of mouse MEIS1.	H, M	ChIP, EMSA, IHC, IP, WB	100 µg 10 µg	39795 39796
NEW Menin antibody (pAb)	Raised against a peptide to the C-terminal region of human Menin.	H	WB	100 µl 10 µl	61005 61006
MEP50 antibody (pAb)	Raised against His-tagged, full-length human MEP50.	H, WR	WB	200 µl 10 µl	39190 39191
MGMT antibody (pAb)	Raised against a peptide to the C-terminal region of human MGMT.	H	WB	100 µl 10 µl	39951 39952
Mi-2 beta antibody (mAb) (Clone 3F2/4)	Raised against a peptide corresponding to human Mi-2 beta.	H, M	IP, WB	100 µg 10 µg	39695 39696
Mi-2 beta antibody (pAb)	Raised against amino acids 22-155 of recombinant mouse Mi-2 beta.	H, M	IF, WB	200 µl 10 µl	39289 39290
MITF antibody (mAb)	Raised against an N-terminal fragment of human MITF.	H	ChIP, EMSA, IF, IHC, IP, WB	100 µg 10 µg	39789 39790
MLH-1 antibody (mAb)	Raised against a peptide corresponding to human MLH-1.	H	WB	100 µg	39219
MLL1/HRX antibody (mAb) (Clone N4.4)	Raised against a recombinant protein containing amino acids 161-356 of mouse MLL1.	H, M	ChIP, IP, WB	100 µg 10 µg	39829 39830
MMSET / WHSC1 antibody (mAb) (Clone 29D1)	Raised against a recombinant protein containing amino acids 1-647 of human MMSET-1.	H	ChIP, IF, IP, WB	100 µg 10 µg	39879 39880
MOZ antibody (pAb)	Raised against a peptide within the N-terminal half of MOZ.	H	WB	100 µl 10 µl	39867 39868
MPP8 antibody (mAb) (Clone N119/44)	Raised against a recombinant protein containing amino acids 50-120 of human MPP8.	H	ICC, WB	100 µg 10 µg	39819 39820
MRE11 antibody (pAb)	Raised against a peptide corresponding to human MRE11.	H	WB	100 µg	39220
MRG15 antibody (pAb)	Raised against a peptide to the C-terminus of human MRG15.	H, WR	WB	200 µl 10 µl	39361 39362
MTA2 antibody (pAb)	Raised against a peptide to the C-terminus of human MTA2.	H, M, WR	WB	200 µl 10 µl	39359 39360

Reactivity: B = Bovine; BY = Budding Yeast; Ce = *C. elegans*; Dr = *Drosophila*; FY = Fission Yeast; Hm = Hamster; H = Human; Mk = Monkey; M = Mouse; NS = Not Species Specific; Rb = Rabbit; R = Rat; WR = Wide range predicted
Applications: ChIP = Chromatin IP; DB = Dot Blot; FC = Flow Cytometry; ICC = Immunocytochemistry; IF = Immunofluorescence; IHC = Immunohistochemistry; IP = Immunoprecipitation; MeDIP = Methyl-DNA IP; WB = Western Blot

Antibody	Immunogen	Reactivity	Applications	Format	Catalog No.
Myc Tag antibody (mAb)	Raised against a peptide to amino acids 410-419 of human c-Myc.	WR	WB	200 µg 10 µg	39279 39280
MyD88 antibody (pAb)	Raised against amino acids 233-248 of human MyD88.	H	WB	100 µg	40914
Myf-5 antibody (mAb) (Clone 1E2G4)	Raised against a peptide derived from mouse Myf-5.	H	IF, WB	100 µg 10 µg	39801 39802
MyoD antibody (mAb) (Clone 5F11)	Raised against full-length, recombinant mouse MyoD protein.	H, M	IF, WB	100 µg 10 µg	39991 39992
NAK1/Nur77 antibody (pAb)	Raised against amino acids 251-266 of human NAK1/Nur77.	H, M	WB	200 µl	40982
Nap1 antibody (pAb)	Raised against full-length, His-tagged <i>Drosophila</i> Nap1 protein.	Dr	IP, WB	100 µl	39577
NAPIL1 antibody (pAb)	Raised against a peptide within the N-terminal region of NAPIL1.	H	WB	100 µl 10 µl	39841 39842
NBS1 antibody (pAb)	Raised against a peptide to the C-terminus of human NBS1.	H	WB	200 µl 10 µl	39373 39374
NcoA4 / ELE1 antibody (pAb)	Raised against a peptide to the C-terminal region of human NCoA4.	H	WB	100 µl 10 µl	39895 39896
NF-1A antibody (pAb)	Raised against amino acids 478-492 of human NF-1A.	H	WB	200 µl 10 µl	39397 39398
NF-1A antibody (pAb)	Raised against amino acids 478-492 of human NF-1A.	H, R	WB	100 µl	39036
NF-1A antibody (pAb)	Raised against amino acids 478-492 of human NF-1A.	H, R	EMSA	100 µl	39329
NF-1B2 antibody (pAb)	Raised against amino acids 402-415 of human NF-1B2.	H, R	WB	100 µg	39091
NFATc1 antibody (pAb)	Raised against amino acids 911-925 of human NFATc1.	H	WB	100 µg	39084
NFκB p50 antibody (pAb)	Raised against amino acids 363-376 of human NFκB p50.	H, M	WB	100 µl	39037
NFκB p50 antibody (pAb)	Raised against amino acids 363-376 of human NFκB p50.	H, M, R	EMSA	17 rxns	39330
NFκB p65 antibody (mAb)	Raised against amino acids 526-539 of human NFκB p65.	H, M	FC, WB	100 µg	40916
NFκB p65 antibody (pAb)	Raised against a peptide immunogen to N-terminal NFκB p65.	H	WB	200 µl 10 µl	39283 39284
NFκB p65 antibody (pAb)	Raised against a peptide to the C-terminus of human NFκB p65.	H	WB	200 µl 10 µl	39369 39370
NFκB p65 antibody (pAb)	Raised against amino acids 501-515 of human NFκB p65.	H, R	FC, IHC, WB	100 µl	39039
NFκB p65 phospho Ser529 antibody (pAb)	Raised against a peptide containing phospho-serine 529 of human NFκB p65.	H	WB	100 µl 10 µl	39691 39692
NFκB p65 phospho Ser536 antibody (pAb)	Raised against a peptide containing phospho-serine 536 of human NFκB p65.	H	WB	100 µl 10 µl	39675 39676
NFκB p100 antibody (pAb)	Raised against a peptide to the C-terminus of human NFκB p100.	H	WB	100 µl 10 µl	39687 39688
Nucleolin antibody (mAb) (Clone 3G4B20)	Raised against a salt-extracted nuclear matrix preparation from MDCK cells.	H, M, WR	IF, WB	200 µl 10 µl	39541 39542

Reactivity: B = Bovine; BY = Budding Yeast; Ce = *C. elegans*; Dr = *Drosophila*; FY = Fission Yeast; Hm = Hamster; H = Human; Mk = Monkey; M = Mouse; NS = Not Species Specific; Rb = Rabbit; R = Rat; WR = Wide range predicted
Applications: ChIP = Chromatin IP; DB = Dot Blot; FC = Flow Cytometry; ICC = Immunocytochemistry; IF = Immunofluorescence; IHC = Immunohistochemistry; IP = Immunoprecipitation; MeDIP = Methyl-DNA IP; WB = Western Blot

Antibody	Immunogen	Reactivity	Applications	Format	Catalog No.
Nucleophosmin antibody (pAb)	Raised against a peptide within the C-terminal region of human nucleophosmin.	H	WB	100 µl 10 µl	39849 39850
Oct-2 antibody (mAb)	Raised against full-length human Oct-2.	H	EMSA	17 rxns	39604
Oct-4 antibody (pAb)	Raised against a peptide derived from human Oct-4.	H	WB	100 µl 10 µl	39811 39812
Osteoprotegerin (OPG) antibody (mAb)	Raised against amino acids 20-37 of human Osteoprotegerin (OPG).	H	WB	100 µg	40938
p53 antibody (mAb) (Clone DO1)	Raised against full-length recombinant human p53 protein.	H	IP, WB	200 µg 10 µg	39553 39554
p53 antibody (pAb)	Raised against amino acids 367-381 of human p53.	H, M	IHC, WB	100 µl	39041
p53 antibody (pAb)	Raised against amino acids 367-381 of human p53.	H	ChIP, EMSA	17 rxns	39334
p63 antibody (mAb)	Raised against a peptide corresponding to human p63.	H	WB	100 µg	40974
p63 antibody (pAb)	Raised against a recombinant full-length human p63.	H	ChIP, IF, WB	100 µl 10 µl	39739 39740
p73 antibody (mAb)	Raised against full-length human p73.	H	WB	100 µg	40972
p73ΔN antibody (mAb)	Raised against amino acids 2-13 of human p73ΔN.	H	WB	100 µg	40973
p107 antibody (mAb)	Raised against full-length human p107.	H, M	EMSA	17 rxns	39605
PARP-1 N-terminal antibody (pAb)	Raised against the His-tagged, N-terminal half of human PARP-1.	H	ChIP, IHC, IP, WB	200 µl 10 µl	39559 39560
PARP-1 C-terminal antibody (pAb)	Raised against the His-tagged, C-terminal half of human PARP-1.	H	ChIP, IHC, IP, WB	200 µl 10 µl	39561 39562
PARP-2 antibody (pAb)	Raised against a recombinant C-terminal region of human PARP-2.	H	ChIP, IP, WB	100 µl 10 µl	39743 39744
Pax-5 antibody (pAb)	Raised against amino acids 1-15 of human Pax-5.	H	WB	100 µl	39043
Pax-7 antibody (mAb) (Clone 2F12H4)	Raised against a peptide derived from mouse Pax-7.	H	IF, WB	100 µg 10 µg	39803 39804
NEW PAX9 antibody (mAb) (Clone 7C2)	Raised against recombinant protein with amino acids 263-342 of mouse PAX9.	H	IHC, IP, WB	100 µg 10 µg	61077 61078
NEW PCNA antibody (mAb) (Clone 16D10)	Raised against full-length recombinant human PCNA.	H	IF, IP, WB	100 µg 10 µg	61079 61080
PGC1-β antibody (pAb)	Raised against a peptide within the N-terminal region of human PGC1-β.	H	WB	100 µl 10 µl	39859 39860
Phc1 antibody (mAb) (Clone 6-1-3)	Raised against full-length, recombinant mouse Phc1.	H	IF, IP, WB	100 µg 10 µg	39723 39724
Phc2 antibody (mAb)	Raised against a recombinant fragment of human Phc2.	H, M	ChIP, IF, IP	100 µg 10 µg	39661 39662
PHF8 antibody (pAb)	Raised against a peptide derived from human PHF8.	H	WB	100 µl 10 µl	39711 39712

Reactivity: B = Bovine; BY = Budding Yeast; Ce = *C. elegans*; Dr = *Drosophila*; FY = Fission Yeast; Hm = Hamster; H = Human; Mk = Monkey; M = Mouse; NS = Not Species Specific; Rb = Rabbit; R = Rat; WR = Wide range predicted
Applications: ChIP = Chromatin IP; DB = Dot Blot; FC = Flow Cytometry; ICC = Immunocytochemistry; IF = Immunofluorescence; IHC = Immunohistochemistry; IP = Immunoprecipitation; MeDIP = Methyl-DNA IP; WB = Western Blot

Antibody	Immunogen	Reactivity	Applications	Format	Catalog No.
PLZF antibody (mAb) (Clone 2A9)	Raised against full-length, recombinant mouse PLZF protein with a GST tag.	H	ChIP, IF, WB	100 µg 10 µg	39987 39988
PML antibody (mAb) (Clone 36.1-104)	Raised against amino acids 1-581 of mouse PML.	M	IF, IP, WB	100 µg 10 µg	39773 39774
PMS2 antibody (mAb)	Raised against amino acids 623-639 of human PMS2.	H, M	WB	100 µg	39221
PP2A antibody (pAb)	Raised against amino acids 288-303 of human PP2A.	H, M, R, WR	ChIP, IP, WB	200 µl 10 µl	39192 39193
NEW PPAR-γ antibody (mAb) (Clone 3PP-IH4)	Raised against a peptide within the N-terminal region of human PPAR-γ1 and γ2.	H	IF, WB	100 µg 10 µg	61025 61026
NEW Progesterone Receptor antibody (mAb) (Clone 6A)	Raised against a peptide to the C-terminus of human Progesterone Receptor.	H	IF, IHC, WB	100 µg 10 µg	61023 61024
NEW PRMT5 antibody (pAb)	Raised against a peptide within the N-terminal region of human PRMT5.	H	WB	100 µl 10 µl	61001 61002
NEW PRMT6 antibody (pAb)	Raised against a peptide within the N-terminal region of human PRMT6.	H	WB	100 µl 10 µl	61003 61004
Progerin antibody (mAb) (Clone 13A4)	Raised against a peptide with amino acids 604-611 of human Progerin.	H	IF, IP, WB	100 µg 10 µg	39965 39966
NEW Prox1 antibody (mAb) (Clone 4G10)	Raised against a recombinant protein with amino acids 547-737 of human Prox1.	H	WB	100 µg 10 µg	61091 61092
NEW Prox1 antibody (mAb) (Clone 5G10)	Raised against a recombinant protein with amino acids 547-737 of human Prox1.	H	IF, IHC	100 µg 10 µg	61093 61094
PXR antibody (pAb)	Raised against a peptide with amino acids 51-71 of human PXR.	H	WB	200 µl 10 µl	39589 39590
Rad1 antibody (pAb)	Raised against amino acids 133-148 of human Rad1.	H	WB	100 µg	40978
Rad17 phospho Ser647 antibody (pAb)	Raised against a peptide with phospho-serine 647 of mouse Rad17.	H, R	IF, WB	100 µg	39222
Rad18 antibody (mAb)	Raised against amino acids 402-414 of human Rad18.	H	WB	100 µg	40979
Rad21 antibody (mAb) (Clone 52A311)	Recombinant protein from the N-terminal half of mouse Rad21.	H, M	WB	200 µg 10 µg	39383 39384
Rad51 antibody (pAb)	Raised against full-length human Rad51 expressed in <i>E. coli</i> .	H	IF, IP, WB	200 µg 10 µg	39194 39195
Rad52 antibody (pAb)	Raised against full-length human Rad52 expressed in <i>E. coli</i> .	H	IF, IHC	200 µg 10 µg	39196 39197
RANK antibody (mAb)	Raised against amino acids 326-616 of GST-tagged, human RANK.	H	WB	100 µg	40917
NEW RAP1 / TERF2IP antibody (pAb)	Raised against a peptide within human RAP1 / TERF2IP.	H	WB	100 µl 10 µl	61069 61070
RAR-α antibody (mAb) (Clone 9α-9A6)	Raised against recombinant protein with amino acids 420-443 of human RAR-α.	H, M	ELISA, EMSA, ICC, IHC, IP, WB	100 µg 10 µg	39971 39972

Reactivity: B = Bovine; BY = Budding Yeast; Ce = *C. elegans*; Dr = *Drosophila*; FY = Fission Yeast; Hm = Hamster; H = Human; Mk = Monkey; M = Mouse; NS = Not Species Specific; Rb = Rabbit; R = Rat; WR = Wide range predicted
Applications: ChIP = Chromatin IP; DB = Dot Blot; FC = Flow Cytometry; ICC = Immunocytochemistry; IF = Immunofluorescence; IHC = Immunohistochemistry; IP = Immunoprecipitation; MeDIP = Methyl-DNA IP; WB = Western Blot

	Antibody	Immunogen	Reactivity	Applications	Format	Catalog No.
NEW	RAR- β antibody (mAb) (Clone 8B-10B2)	Raised against amino acids 413-448 of human RAR- β .	H	ELISA, IHC, IP, WB	100 μ g 10 μ g	61027 61028
	RAR- β 2 antibody (pAb)	Raised against amino acids 36-53 of human RAR- β 2.	H	WB	100 μ l	39050
	RAR- γ antibody (pAb)	Raised against amino acids 412-428 of mouse RAR- γ .	H, M	WB	100 μ l	39051
	RAR- γ antibody (mAb) (Clone 4 γ -7A11)	Raised against recombinant protein with amino acids 428-441 of human RAR- γ .	H	ELISA, EMSA, ICC, IP, WB	100 μ g 10 μ g	39973 39974
	Rb antibody (mAb) (Clone IF8)	Raised against full-length human Rb.	H, M	EMSA	17 rxns	39607
	Rb antibody (mAb) (Clone XZ77)	Raised against full-length human Rb.	H	WB	100 μ l	39520
	Rb antibody (mAb) (Clone XZ77)	Raised against full-length human Rb.	H	EMSA	17 rxns	39606
	Rb antibody (pAb)	Raised against amino acids 881-895 of human Rb.	H, M, R	WB	100 μ l	39052
	RbAp46/48 antibody (pAb)	Raised against recombinant human RbAp46 protein.	H, WR	ChIP, WB	200 μ l 10 μ l	39198 39199
	RelB antibody (pAb)	Raised against amino acids 476-490 of human RelB.	H	EMSA	17 rxns	39339
	Ring1B antibody (mAb)	Raised against recombinant full-length human Ring1B.	H, Mk, M	ChIP, IF, IP, WB	100 μ g 10 μ g	39663 39664
NEW	RNA Pol II antibody (mAb) (Clone 4F8)	Raised against peptide with human RNA Pol II C-terminal heptad repeat domain.	H	ChIP, IF, IP, WB	100 μ g 10 μ g	61081 61082
	RNA pol II antibody (mAb)	Raised against a peptide containing YSPSPS from human RNA pol II.	BY, Ce, H, M, R	ChIP, IF, IP, WB	200 μ l	39097
NEW	RNA Pol II CTD phospho Ser2 antibody (mAb) (Clone 3E10)	Peptide with the RNA Pol II CTD sequence phosphorylated at serine 2.	H	ChIP, IF, IP, WB	100 μ g 10 μ g	61083 61084
	RNA Pol II CTD phospho Ser2 antibody (pAb)	Peptide with the RNA Pol II CTD sequence phosphorylated at serine 2.	H	WB	200 μ l 10 μ l	39563 39564
NEW	RNA Pol II CTD phospho Ser5 antibody (mAb) (Clone 3E8)	Peptide with the RNA Pol II CTD sequence phosphorylated at serine 5.	H	ChIP, IF, IP, WB	100 μ g 10 μ g	61085 61086
	RNA pol II CTD phospho Ser5 antibody (pAb)	Peptide with RNA Pol II heptad repeat phosphorylated at serine 5.	H, WR	ChIP, IF, WB	200 μ l 10 μ l	39233 39234
	RNA pol II CTD phospho Ser5 antibody (pAb)	Peptide with RNA Pol II heptad repeat phosphorylated at serine 5.	H, WR	ChIP, WB	100 μ g 10 μ g	39749 39750
NEW	RNA Pol II CTD phospho Ser7 antibody (mAb) (Clone 4E12)	Peptide with the RNA Pol II CTD sequence phosphorylated at serine 7.	H	ChIP, IF, IP, WB	100 μ g 10 μ g	61087 61088
	Rsf1 antibody (pAb)	Raised against His-tagged protein with amino acids 165-350 <i>Drosophila</i> Rsf1.	Dr	IF, IP, WB	100 μ l	39579
	RXR- α antibody (pAb)	Raised against amino acids 14-28 of human RXR- α .	H	WB	100 μ l	39054
NEW	RXR- α antibody (mAb) (Clone 4RX-3A2)	Raised against recombinant protein to the C-terminal region of mouse RXR- α .	H	EMSA, IP, WB	100 μ g 10 μ g	61029 61030
	RXR- β antibody (mAb) (Clone 16RX-2E1)	Raised against peptide to the N-terminal region of mouse RXR- β .	H	EMSA, IF, IHC, WB	100 μ g 10 μ g	61031 61032

Reactivity: B = Bovine; BY = Budding Yeast; Ce = *C. elegans*; Dr = *Drosophila*; FY = Fission Yeast; Hm = Hamster; H = Human; Mk = Monkey; M = Mouse; NS = Not Species Specific; Rb = Rabbit; R = Rat; WR = Wide range predicted
Applications: ChIP = Chromatin IP; DB = Dot Blot; FC = Flow Cytometry; ICC = Immunocytochemistry; IF = Immunofluorescence; IHC = Immunohistochemistry; IP = Immunoprecipitation; MeDIP = Methyl-DNA IP; WB = Western Blot

	Antibody	Immunogen	Reactivity	Applications	Format	Catalog No.
NEW	RXR- γ antibody (mAb) (Clone 12RX-2D3)	Raised against peptide within the D-region region of mouse RXR- γ .	H	EMSA, IHC, WB	100 μ g 10 μ g	61033 61034
	SAF-1 antibody (pAb)	Raised against amino acids 180-193 of human SAF-1.	H	WB	100 μ l	39056
	SALL4 antibody (pAb)	Raised against a peptide to the N-terminus of human SALL4.	H	WB	100 μ l 10 μ l	39957 39958
	SAP30 antibody (pAb)	Raised against recombinant full-length human SAP30.	H	IP, WB	100 μ l 10 μ l	39731 39732
	SATB1 antibody (pAb)	Raised against a peptide within the N-terminal region of SATB1.	H	WB	100 μ l 10 μ l	39839 39840
	NEW	SET8 / PR-SET7 antibody (pAb)	Raised against a peptide to the N-terminal half of SET8 / PR-SET7.	H	WB	100 μ l 10 μ l
SIN3A antibody (pAb)		Raised against a peptide to the N-terminus of SIN3A.	H	WB	100 μ l 10 μ l	39865 39866
SIRT1 antibody (mAb) (Clone 2G1/F7)		Raised against amino acids 581-630 of human SIRT1.	H	IF, IP, WB	200 μ g 10 μ g	39353 39354
SIRT2 antibody (pAb)		Raised against a peptide to the C-terminus of human SIRT2.	H	WB	100 μ l 10 μ l	39903 39904
NEW	SIRT2 antibody (pAb)	Raised against a peptide to the C-terminus of human SIRT2.	H	IF, IP, WB	100 μ l 10 μ l	61043 61044
NEW	SIRT2 antibody (pAb)	Raised against a peptide to the C-terminus of mouse SIRT2.	M	IF, WB	100 μ l 10 μ l	61045 61046
	SIRT6 antibody (pAb)	Raised against a peptide to the C-terminus of human SIRT6.	H	WB	100 μ l 10 μ l	39911 39912
NEW	SMC1- α antibody (pAb)	Raised against a peptide to the C-terminus of human SMC1- α .	H	WB	100 μ l 10 μ l	61067 61068
	SMC1 phospho Ser957 antibody (pAb)	Raised against a peptide with phospho-serine 957 of human SMC1.	H	WB	200 μ l 10 μ l	39631 39632
	SMRT / NCoR2 antibody (pAb)	Raised against a peptide within the N-terminal region of SMRT / NCoR2.	H	WB	100 μ l 10 μ l	39845 39846
	SNF2h antibody (mAb)	SNF2h+ clones were selected from purified RSF complex-immunized mice.	H, M	ChIP, IF, IP, WB	200 μ l 10 μ l	39543 39544
	SNF5 antibody (pAb)	Raised against a peptide to the C-terminus of human SNF5.	H	WB	100 μ l 10 μ l	39775 39776
	Sox2 antibody (pAb)	Raised against a peptide containing amino acids 111-140 of mouse Sox2.	H	ChIP, IF, IHC, IP, WB	100 μ l 10 μ l	39823 39824
Sox2 antibody (pAb)	Raised against a peptide within the C-terminal region of human Sox2.	H	WB	100 μ l 10 μ l	39843 39844	
Sp1 antibody (pAb)	Raised against amino acids 520-534 of human Sp1.	H, M, R	WB	100 μ l	39058	

Reactivity: B = Bovine; BY = Budding Yeast; Ce = *C. elegans*; Dr = *Drosophila*; FY = Fission Yeast; Hm = Hamster; H = Human; Mk = Monkey; M = Mouse; NS = Not Species Specific; Rb = Rabbit; R = Rat; WR = Wide range predicted
Applications: ChIP = Chromatin IP; DB = Dot Blot; FC = Flow Cytometry; ICC = Immunocytochemistry; IF = Immunofluorescence; IHC = Immunohistochemistry; IP = Immunoprecipitation; MeDIP = Methyl-DNA IP; WB = Western Blot

Antibody	Immunogen	Reactivity	Applications	Format	Catalog No.
Sp1 non-phospho antibody (pAb)	Raised against a peptide with serine 101 of human Sp1. This antibody recognizes human Sp1 when it is not phosphorylated at serine 101.	H	WB	100 µl	39809
				10 µl	39810
Sp1 phospho Ser101 antibody (pAb)	Raised against a peptide with phospho-serine 101 of human Sp1.	H	WB	100 µl	39757
				10 µl	39758
SREBP-1 antibody (mAb) (Clone IgG-2A4)	Raised against a recombinant protein with amino acids 301-407 of human SREBP-1.	H	IP, WB	100 µg	39939
				10 µg	39940
SREBP-2 antibody (mAb) (Clone IgG-1C6)	Raised against a recombinant protein with amino acids 833-1141 of human SREBP-2.	H	ChIP, IP, WB	100 µg	39941
				10 µg	39942
SRF Drosophila antibody (mAb)	Raised against full-length Drosophila SRF.	Dr	WB	200 µl	39093
STAT1 phospho Ser727 antibody (pAb)	Raised against a peptide with phospho-serine 727 of human STAT1.	H	WB	200 µl	39633
				10 µl	39634
STAT1α antibody (pAb)	Raised against amino acids 712-750 of human STAT1α.	H	WB	100 µl	39059
STAT2 antibody (pAb)	Raised against amino acids 830-849 of human STAT2.	H, M	WB	100 µg	39060
STAT2 phospho Tyr689 antibody (pAb)	Raised against a peptide with phospho-tyrosine 689 of human STAT2.	H	WB	200 µl	39611
				10 µl	39612
STAT3 phospho Tyr705 antibody (pAb)	Raised against a peptide with phospho-tyrosine 705 of human STAT3.	H	WB	200 µl	39595
				10 µl	39596
STAT3 phospho Ser727 antibody (pAb)	Raised against a peptide with phospho-tyrosine 727 of human STAT3.	H	WB	200 µl	39613
				10 µl	39614
STAT4 antibody (pAb)	Raised against amino acids 731-748 of mouse STAT4.	H, M, R	WB	100 µl	39062
STAT5A/B phospho Tyr694/Tyr699 antibody (pAb)	Raised against a peptide with phospho-tryosine 694 of human STAT5A. (Human STAT5B shares the identical sequence surrounding Ser731.)	H	WB	200 µl	39617
				10 µl	39618
STAT5A/B phospho Ser726/Ser731 antibody (pAb)	Raised against a peptide with phospho-serine 726 of human STAT5A. (Human STAT5B shares the identical sequence surrounding Ser731.)	H	WB	200 µl	39597
				10 µl	39598
STAT5B antibody (pAb)	Raised against amino acids 777-787 of human STAT5B.	H	WB	100 µl	39064
SUV39H1 antibody (mAb) (Clone MG44)	Raised against a recombinant protein with amino acids 1-126 of mouse SUV39H1.	H, M	ChIP, IP, WB	100 µg	39785
				10 µg	39786
Suz12 antibody (mAb) (Clone 2A09)	Raised against full-length recombinant human Suz12.	H, WR	ChIP, IF, IHC, IP	100 µg	39877
				10 µg	39878
Suz12 antibody (pAb)	Raised against a peptide to the C-terminus of human Suz12.	H, WR	WB	200 µl	39357
				10 µl	39358
TAC1 antibody (pAb)	Raised against amino acids 116-132 of human TAC1.	H, M	FC, WB	100 µg	40936
TBK1 antibody (mAb)	Raised against amino acids 563-577 of human TBK1.	H, M, R	WB	100 µg	40909
TLR2 antibody (pAb)	Peptide mix of amino acids 180-196, 353-370 & 473-489 of human TLR2.	H	WB	100 µg	40981
TLR3 antibody (mAb)	Raised against a KLH-peptide to human TLR3.	H	FC, IHC, IP, WB	100 µg	40952
TLR6 antibody (mAb)	Raised against amino acids 408-424 of human TLR6.	H	FC, WB	100 µg	40954
TLR8 antibody (mAb)	Raised against amino acids 750-850 of human TLR8.	H, M	FC, WB	100 µg	40955

Reactivity: B = Bovine; BY = Budding Yeast; Ce = *C. elegans*; Dr = *Drosophila*; FY = Fission Yeast; Hm = Hamster; H = Human; Mk = Monkey; M = Mouse; NS = Not Species Specific; Rb = Rabbit; R = Rat; WR = Wide range predicted
Applications: ChIP = Chromatin IP; DB = Dot Blot; FC = Flow Cytometry; ICC = Immunocytochemistry; IF = Immunofluorescence; IHC = Immunohistochemistry; IP = Immunoprecipitation; MeDIP = Methyl-DNA IP; WB = Western Blot

Antibody	Immunogen	Reactivity	Applications	Format	Catalog No.
TLR9 antibody (mAb)	Raised against a KLH-peptide to amino acids 268-284 of human TLR9.	H, M, R	FC, IHC, WB	100 µg	40956
TORC2 antibody (pAb)	Raised against a peptide to the C-terminus of human TORC2.	H, M	WB	100 µl 10 µl	39779 39780
TRAF2 antibody (mAb)	Raised against amino acids 215-222 of human TRAF2 protein.	H	WB	100 µg	40919
TRAF5 antibody (mAb)	Raised against amino acids 77-186 of human TRAF5 protein.	H	WB	100 µg	40920
TRAIL antibody (mAb)	Raised against amino acids 17-35 of human TRAIL.	H	FC, WB	100 µg	40966
TRANCE/RANKL antibody (mAb)	Raised against amino acids 1-317 of GST-tagged, human TRANCE/RANKL.	H, M	FC, IHC, WB	100 µg	40940
TRF2 antibody (pAb)	Raised against full-length human TRF2.	H, M, R	ChIP, IP, WB	100 µg	39223
TROY/TAJ antibody (pAb)	Peptide mix of amino acids 29-44, 204-219 & 409-421 of human TROY/TAJ.	H, M	WB	100 µg	40937
NEW TWIST antibody (mAb)	Raised against full-length recombinant mouse TWIST protein.	H	IF, IHC, WB	100 µg 10 µg	61097 61098
UBE2N antibody (pAb)	Peptide mixture of amino acids 2-19 and 131-148 of human UBE2N.	H	IP, WB	100 µg	39224
UBE2V1α antibody (pAb)	Raised against amino acids 31-49 of human UBE2V1α.	H	WB	100 µg	39225
UBE2V2 antibody (pAb)	Raised against amino acids 12-28 of human UBE2V2.	H	WB	100 µg	39226
Ubiquitin antibody (mAb) (Clone P4D1)	Raised against purified bovine ubiquitin.	WR	IF, IHC, IP, WB	100 µg 10 µg	39741 39742
Ubn1 antibody (pAb)	Raised against amino acids 1-345 of human Ubn1.	H	WB	100 µl 10 µl	39869 39870
Uhrf1 antibody (pAb)	Raised against a peptide to the C-terminus of human Uhrf1.	H	WB	200 µl 10 µl	39625 39626
USF1 antibody (pAb)	Raised against amino acids 1-14 of human USF1.	H	WB	100 µl	39068
USF1 antibody (pAb)	Raised against amino acids 1-14 of human USF1.	H, R	EMSA	17 rxns	39343
YY1 antibody (pAb)	Raised against amino acids 109-123 of human YY1.	H, Mk, M, R	WB	100 µl	39071
YY1 antibody (pAb)	Raised against amino acids 109-123 of human YY1.	H	EMSA	17 rxns	39345

Reactivity: B = Bovine; BY = Budding Yeast; Ce = *C. elegans*; Dr = *Drosophila*; FY = Fission Yeast; Hm = Hamster; H = Human; Mk = Monkey; M = Mouse; NS = Not Species Specific; Rb = Rabbit; R = Rat; WR = Wide range predicted
Applications: ChIP = Chromatin IP; DB = Dot Blot; FC = Flow Cytometry; ICC = Immunocytochemistry; IF = Immunofluorescence; IHC = Immunohistochemistry; IP = Immunoprecipitation; MeDIP = Methyl-DNA IP; WB = Western Blot

Fluorescent and HRP Secondary Antibody Conjugates

The table below provides details on all of Active Motif's secondary antibody conjugates. For more comprehensive information on each, including data and downloadable Technical Data Sheets, please visit our website at www.activemotif.com/secondary.

Name	Absorption	Emission	ϵ L/(mol-cm)	Stokes shift	Format	Catalog No.
HRP Goat anti-Mouse IgG	—	—	—	—	400 μ g	15014
HRP Goat anti-Rabbit IgG	—	—	—	—	400 μ g	15015
ATTO 594 Goat anti-Mouse IgG	601 nm	627 nm	120,000	26 nm	250 μ l 35 μ l	15037 15057
ATTO 594 Goat anti-Rabbit IgG	601 nm	627 nm	120,000	26 nm	250 μ l 35 μ l	15047 15067
ATTO 647N (STED) Goat anti-Mouse IgG	644 nm	669 nm	150,000	25 nm	250 μ l 35 μ l	15038 15058
ATTO 647N (STED) Goat anti-Rabbit IgG	644 nm	669 nm	150,000	25 nm	250 μ l 35 μ l	15048 15068
ATTO 655 (STED) Goat anti-Mouse IgG	663 nm	684 nm	125,000	21 nm	250 μ l 35 μ l	15039 15059
ATTO 655 (STED) Goat anti-Rabbit IgG	663 nm	684 nm	125,000	21 nm	250 μ l 35 μ l	15049 15069
Chromo™ 488 Goat anti-Mouse IgG	488 nm	517 nm	73,000	29 nm	1 mg 100 μ g	15031 15051
Chromo™ 488 Goat anti-Rabbit IgG	488 nm	517 nm	73,000	29 nm	1 mg 100 μ g	15041 15061
Chromo™ 494 Goat anti-Mouse IgG	494 nm	628 nm	55,000	124 nm	1 mg 100 μ g	15032 15052
Chromo™ 494 Goat anti-Rabbit IgG	494 nm	628 nm	55,000	124 nm	1 mg 100 μ g	15042 15062
Chromo™ 505 Goat anti-Mouse IgG	505 nm	526 nm	70,000	21 nm	1 mg 100 μ g	15030 15050
Chromo™ 505 Goat anti-Rabbit IgG	505 nm	526 nm	70,000	21 nm	1 mg 100 μ g	15040 15060
Chromo™ 546 Goat anti-Mouse IgG	545 nm	561 nm	98,800	16 nm	1 mg 100 μ g	15033 15053
Chromo™ 546 Goat anti-Rabbit IgG	545 nm	561 nm	98,800	16 nm	1 mg 100 μ g	15043 15063
Chromo™ 642 Goat anti-Mouse IgG	642 nm	660 nm	180,000	18 nm	1 mg 100 μ g	15034 15054
Chromo™ 642 Goat anti-Rabbit IgG	642 nm	660 nm	180,000	18 nm	1 mg 100 μ g	15044 15064